



# The Open Orthopaedics Journal

Content list available at: [www.benthamopen.com/TOORTHJ/](http://www.benthamopen.com/TOORTHJ/)

DOI: 10.2174/1874325001610010481



## RESEARCH ARTICLE

# Comparing Transtibial and Anteromedial Drilling Techniques for Single-bundle Anterior Cruciate Ligament Reconstruction

Erhan Sukur<sup>a,\*</sup>, Yunus Emre Akman<sup>b</sup>, Ahmet Senel<sup>c</sup>, Ethem Ayhan Unkar<sup>d</sup>, Huseyin Nevzat Topcu<sup>a</sup>, and Yusuf Ozturkmen<sup>c</sup>

<sup>a</sup>Sakarya Education and Research Hospital, 54100, Sakarya, Turkey

<sup>b</sup>Metin Sabanci Baltalimani Bone and Diseases Training and Research Hospital, 34470, Istanbul, Turkey

<sup>c</sup>Istanbul Education and Research Hospital, 34100, Istanbul, Turkey

<sup>d</sup>Kartal Yavuz Selim State Hospital, 34050, Istanbul, Turkey

Received: July 09, 2016

Revised: August 01, 2016

Accepted: September 11, 2016

### Abstract:

#### Background:

Among the many factors that determine the outcome following anterior cruciate ligament (ACL) reconstruction, the position of the femoral tunnel is known to be critically important and is still the subject of extensive research.

#### Objective:

We aimed to retrospectively compare the outcomes of arthroscopic ACL reconstruction using transtibial (TT) or anteromedial (AMP) drilling techniques for femoral tunnel placement.

#### Methods:

ACL reconstruction was performed using the TT technique in 49 patients and the AMP technique in 56 patients. Lachman and pivot-shift tests, the Lysholm Knee Scale, International Knee Documentation Committee (IKDC) score, Tegner activity scale and visual analog scale (VAS) were used for the clinical and functional evaluation of patients. Time to return to normal life and time to jogging were assessed in addition to the radiological evaluation of femoral tunnel placement.

#### Results:

In terms of the Lysholm, IKDC, Tegner score, and stability tests, no significant differences were found between the two groups ( $p > 0.05$ ). Statistical analysis revealed reduced time to return to normal life and jogging in the AMP group ( $p < 0.05$ ). The VAS score was also significantly reduced in the AMP group ( $p < 0.05$ ). The position of the femoral tunnel was anatomically appropriate in 51 patients in the AMP group and 5 patients in the TT group.

#### Conclusion:

The AMP technique is superior to the TT technique in creating anatomical femoral tunnel placement during single-bundle ACL reconstruction and provides faster recovery in terms of return to normal life and jogging at short-term follow-up.

**Keywords:** Anterior cruciate ligament, Anteromedial portal, Drilling, Femoral tunnel, Graft, Transtibial technique.

\* Address correspondence to this author at the Sakarya Education and Research Hospital, 54100, Sakarya, Turkey; Tel: +905322632090; Fax: +902642759192; E-mail: [erhan\\_sukur@hotmail.com](mailto:erhan_sukur@hotmail.com)

## INTRODUCTION

Surgical reconstruction is now widely accepted as the treatment of choice for functional knee instability due to anterior cruciate ligament (ACL) deficiency especially in young patients who are involved in physically high demanding activities [1]. This procedure is well-known for its ability to allow an individual to return to preinjury activity levels, which may not be achieved with non-surgical treatment [2]. In spite of the considerable research focused on ACL reconstruction, the increased risk of early knee osteoarthritis and the presence of rotational instability post-surgery have not been fully investigated yet. As a result, controversy remains regarding the best technique for reconstruction; thus, it is still the subject of extensive research [3, 4]. Among the many factors that determine the outcome following ACL reconstruction, the position of the tunnels created for graft fixation is known to be of critical influence, and malpositioning of the femoral tunnel occurs 3 times more frequently than malpositioning of the tibial tunnel [3, 5, 6]. Femoral tunnel position is more critical because it is closer to the central axis of knee rotation, and malpositioning of the femoral tunnel has been reported to be the most frequent cause of graft failure [7, 8]. A non-anatomically placed graft can create abnormal tensile and compressive forces, which affect graft healing and re-integration processes [9].

Anatomical reconstruction of the ACL involves the placement of the femoral tunnel at the center of the original femoral footprint [10]. Recent studies have shown that anatomic positioning of the femoral tunnel best restores native knee kinematics by allowing appropriate graft tension throughout the range of motion, and results show increased rotational stability, as well as translational and tensioning patterns similar to the native ACL [11, 12]. The transtibial (TT) technique is the most widely used method for drilling the femoral tunnel during ACL reconstruction, but positioning the femoral tunnel anatomically is difficult and constrained by the position of the tibial tunnel. As a result, the graft is oriented in a relatively vertical direction, in the footprint of the anteromedial bundle of the ACL [13]. Thus, over the past decade, there has been a trend transitioning from use of the TT drilling technique toward the anteromedial portal (AMP) drilling technique in order to achieve more anatomical femoral tunnel placement. The AMP technique requires independent drilling through an additional anteromedial portal in a more horizontal direction [3, 14] (Fig. 1).



**Fig. (1).** Frontal magnetic resonance imaging (MRI) demonstrating the horizontal orientation of the graft in the femoral tunnel.

According to anatomical studies, the AMP technique currently allows for a more anatomical femoral tunnel position within the native ACL femoral footprint compared with the TT technique. However, no adequate studies exist that

compare the clinical and functional outcomes of these two techniques [2, 9]. Thus, this study aimed to retrospectively compare the radiological and clinical outcomes of arthroscopic ACL reconstruction using the TT and AMP drilling techniques for femoral tunnel placement.

## MATERIALS AND METHODS

### Patients

Among the patients who underwent single-bundle ACL reconstruction between February 2011 and July 2014 using an autologous hamstring graft, 105 eligible patients were included in this retrospective study. ACL reconstruction was performed using the TT technique in 49 patients (4 female, 45 male; mean age, 26.8 years, range 18-40) and the AMP technique in 56 patients (4 female, 52 male; mean age, 25.5 years, range 17-35). The inclusion criteria for this study were as follows: less than 6 months between injury and surgery, at least 24 months of follow-up, a healthy contralateral knee, and patient age between 16 and 40 years. Patients who underwent concomitant surgery during ACL reconstruction for other combined injuries, such as collateral ligament injury, posterior cruciate ligament injury, or posterolateral corner injury were excluded from the study. All patients were operated by two senior surgeons. This study was approved by our institutional ethics committee and informed consent was obtained from all patients for the use of their demographic and radiological data.

### Surgical Technique

Arthroscopic examination was performed and ACL rupture was confirmed before harvesting the graft. Quadruple-strand semitendinosus and gracilis autografts were used in all patients. The standard technique was performed to place the tibial tunnel in both the TT and AMP groups [15]. The tibial tunnel was prepared in the footprint of the ACL at an angle of 45° to the tibial shaft. In the TT group, the standard aiming guide with a 7-mm offset was placed through the tibial tunnel at the 11 o'clock position of the right knee and the 1 o'clock position of the left knee, and a pin was advanced to determine the femoral tunnel position. A cannulated reamer was then used transtibially to create the femoral tunnel. In the AMP group, a three-portal technique (anterolateral, central anteromedial, and low anteromedial) was used. The low AMP was created under arthroscopic visualization; a spinal needle was advanced into the joint just above the anterior horn of the medial meniscus and 1.5 cm medial to the medial border of patellar tendon and to create the femoral tunnel independently from the tibial tunnel. It is important to keep a security distance at least 2 mm from the cartilage of the medial femoral condyle once the drill passes through this portal [16]. The midpoints of the remnants of the anteromedial and posterolateral bundles of the ACL were marked with a thermal device. The lateral intercondylar ridge and lateral bifurcate ridge served as the bony landmarks for the femoral attachments. The lateral intercondylar ridge is an osseous landmark in the medial wall of the femoral lateral condyle that runs from proximal to distal and anterior to posterior in the anatomic position. There is no ACL attachment anterior to this ridge. Lateral bifurcate ridge is an osseous landmark that runs from anterior to posterior and separates the AM and PL bundles femoral attachment site [17]. The knee was flexed to 120°-130°, and the guidewire was placed in the center of the two insertion areas via the low AMP. Femoral drills were selected according to the graft diameter, and the tunnel was drilled with a cannulated reamer using a freehand technique. An endobutton continuous loop device was used for femoral fixation of the graft. Pre-tensioning of the graft was performed by flexing and extending the knee through the range of motion. The absence of graft impingement was confirmed by arthroscopic examination. Tibial fixation was performed in 20° of flexion using a bioabsorbable screw and a U staple with a posterior drawer.

### Postoperative Care

The same postoperative accelerated rehabilitation protocols were used for both groups. Knee braces or sleeves were not employed in any of the patients. All patients were allowed weight-bearing as tolerated with crutches on the day of surgery. After three weeks, weight-bearing without crutches was permitted.

### Clinical and Radiological Assessment

At the last follow-up, all patients were examined by an orthopedic surgeon who was blinded to the relevant study details and who was not the operating surgeon. Lachman and pivot-shift tests were performed to evaluate the anterior-posterior and rotational stability, respectively. The Lysholm Knee Scale, International Knee Documentation Committee (IKDC) scoring system, Tegner activity level, and visual analog scale (VAS) for patient satisfaction were used for the clinical and functional evaluation of patients before surgery and at the last follow-up [18, 19]. The time to return to

normal life and time to return to jogging were assessed. For the radiological evaluation of tunnel placement, in order to determine whether the reconstructed ACL fell outside of the anatomical range, the criteria developed by Illingworth *et al.* were used [20].

### Statistical Analysis

Statistical analyses were conducted using SPSS version 15.0 for Windows (SPSS Inc., Chicago, IL). When the numerical values for the two independent groups were normally distributed, Student's *t* test was used, and when normal distribution was not achieved, a Mann-Whitney U test was performed to compare the groups. For categorical variables, the ratios between the groups were analyzed by chi-square test. As the conditions for parametric tests were not met, a Monte Carlo simulation was used to evaluate the numerical variables. A *p* value < 0.05 was considered statistically significant.

### RESULTS

Statistically significant differences were not found between the groups with regard to age, sex, body mass index (BMI), mean follow-up period, and time from injury to surgery (*p* > 0.05) (Table 1). The mean follow-up periods for the AMP and TT groups were 30.5 months (range, 24-42 months) and 28.6 months (range, 26-38 months), respectively. The mechanism of injury involved sports activities in 32 patients (65%) in the TT group and 41 patients (73%) in the AMP group. Assessment of the preoperative and postoperative IKDC and Lysholm scores showed a significant improvement in both groups (*p* < 0.05). In terms of the Lysholm and IKDC scores and Tegner activity levels, no significant differences existed between the groups at the last follow-up evaluation (*p* > 0.05) (Table 2).

**Table 1. Patient demographics.**

	TT Group	AMP Group	p Value
Patients (n)	56	49	
Sex (male: female, n)	52:4	45:4	1
Age (mean ± SD, years)	25.5 ± 5	26.8 ± 5.7	0.406
BMI (mean ± SD, kg/m <sup>2</sup> )	24.9 ± 2.9	25.1 ± 2.7	0.735
Follow-up (mean ± SD, months)	30.1 ± 5.4	30.9 ± 5.6	0.441
Time from injury to surgery (mean ± SD, weeks)	10.3 ± 5.1	10.6 ± 4.7	0.415

**Table 2. Functional evaluation of the groups at the last follow-up.**

Parameters	Anteromedial	Transtibial	p Value
IKDC score	93.1	91.4	0.614
Lysholm score	95.4	93.9	0.545
Tegner activity level	6.3	6.1	0.415

Lachman and pivot-shift tests showed no significant differences between the groups (*p* > 0.05) (Table 3). Statistical analysis revealed that a reduced time was needed to return to normal life and jogging in the AMP group than in the TT group (*p* < 0.05), and a statistically significant difference was observed between the two groups (*p* < 0.001). Moreover, higher patient satisfaction was achieved for the AMP group following surgery according to analysis of VAS scores (*p* < 0.05) (Table 4). AMP technique was found to be superior to the TT technique in creating femoral tunnels within anatomical range, according to the radiological evaluation method of Illingworth (*p* < 0.001). The reconstructed ACL fell within an anatomical range in 51 patients (91%) in the AMP group and 5 patients (10.2%) in the TT group (Fig. 2).

**Table 3. Physical examination.**

Parameters	Anteromedial n (%)	Transtibial n (%)	p Value
Pivot-shift test			0.056
0	42 (75.0)	29 (59.2)	
1	14 (25.0)	19 (38.8)	
2	0	1 (2)	
3	0	0	
Lachman test			0.059

(Table 3) contd.....

Parameters	Anteromedial n (%)	Transtibial n (%)	p Value
0	42 (75.0)	29 (59.2)	
1	14 (25.0)	17 (34.7)	
2	0	3 (6.1)	
3	0	0	

Table 4. VAS scores, time to normal life, and time to jogging for the AMP and TT groups.

	Anteromedial	Transtibial	p Value
VAS score (mean ± SD)	9.4 ± 0.8	8.9 ± 0.8	0.002
Time to normal life (mean ± SD, weeks)	7.5 ± 0.9 (8)	9.1 ± 1.3 (9)	< 0.001
Time to jogging (mean ± SD, weeks)	12.4 ± 1.6 (12)	14.4 ± 2.2 (14)	< 0.001




Fig. (2). The femoral tunnel angle (A) and inclination angle (B) suggest that the reconstructed ACL is within the anatomical range.

**DISCUSSION**

The principal finding of this comparative study is that the AMP technique is superior to the TT technique in achieving anatomical femoral tunnel placement. The patients in the AMP group began jogging and returned to normal life significantly earlier at short-term follow-up compared with patients in the TT group.

The most frequent technical flaw resulting in the failure of ACL reconstruction is the non-anatomical placement of the femoral tunnel [5 - 11]. In a cadaveric study by Gadikota *et al.*, improved outcome after ACL reconstruction was notably related to the maximum femoral ACL footprint coverage achieved by the surgical tunnel [21]. Creation of a femoral tunnel with the TT technique requires a relatively short operative time and has been commonly performed with successful results for a long time [22]. However, when using the TT technique, the femoral tunnel is typically placed anteriorly and higher in the intercondylar notch compared with the original ACL femoral footprint [23, 24]. Only the anteromedial bundle of the ACL may be covered by the TT drilling technique, which results in better restoration of sagittal plane stability than rotational stability, which may remain inadequate [25]. The abnormal femoral tunnel placement observed in the TT technique is attributed to the dependency of the femoral tunnel position on the antecedently drilled tibial tunnel [26]. In an attempt to achieve anatomical femoral tunnel position via the TT technique, Golish suggested a far medial entry into the tibia [27]. Controversially, many authors have concluded that attempting to create an anatomical femoral tunnel position through a medial entry into the tibia would require a starting point too close to the tibial joint line, resulting in a relatively short medial tibial tunnel and a tunnel length-graft length mismatch [23, 26]. Youm *et al.* used a modified TT technique, positioning the tibia in internal rotation and varus alignment while

creating the femoral tunnel. They noted that adjusting the guidewire within the native ACL site using the modified TT technique is easily possible [28].

On the other hand, numerous studies have suggested that the AMP technique facilitates a more anatomic and horizontal femoral tunnel, independent of tibial tunnel placement within the native ACL femoral attachment site, and provides greater rotational stability. Furthermore, these studies suggest that a more horizontal placement of the graft decreases the risk of graft impingement [10, 29, 30]. However, there are several potential surgical difficulties with regard to the AMP technique, including a short femoral tunnel, blow out of the back wall of the lateral femoral intercondylar notch, and difficulty with visualization and graft passage [31]. Nevertheless, in the present study, the AMP technique was found to be superior to the TT technique in creating femoral tunnels within anatomical range, according to the radiological evaluation method of Illingworth. Illingworth confirmed with 3D computed tomography that a femoral tunnel angle greater than  $32.7^\circ$  (100% sensitivity and 85% specificity) on an anteroposterior radiograph of the knee and an inclination angle less than  $55^\circ$  (100% sensitivity and 87.5% specificity) on sagittal magnetic resonance imaging (MRI) determine whether the reconstructed ACL is within the anatomical range [20]. We consider this to be an easy and beneficial method to determine whether the reconstructed ACL is within the anatomical range.

Although evaluation of the functional outcomes based on IKDC and Lysholm scores and Tegner activity levels revealed higher values in the AMP group, significant differences between the groups were not observed. This result is similar to the results of other studies with more than a year of follow-up time [11, 29, 32, 33]. Conversely, there are studies with significantly higher IKDC and/or Lysholm scores following ACL reconstruction using the AMP technique [34]. As a result, no definitive evidence has been found to conclude that one technique is superior to the other in terms of functional outcomes based on scoring systems, such as the IKDC scoring system, Lysholm score, and Tegner activity level, in clinical studies [35].

Knee stability with respect to the ACL can be measured by the amount of anterior tibial translation identified using the Lachman and pivot-shift physical tests or using an arthrometer [34]. Although patients with TT ACL reconstruction obtain good anteroposterior stability, numerous biomechanical studies have demonstrated that the AMP technique restores the translational and rotational stability better and provides more anatomic graft placement [36 - 40]. In this study, anterior translation of the tibia was evaluated using both Lachman and pivot-shift tests, and no statistically significant difference was found between the groups. This finding may be due to the subjective nature of the tests. However, we performed stability tests at the last follow-up, which was at least two years after surgery.

Several clinical and biomechanical studies have demonstrated improved knee stability with the AMP technique, particularly during the first months [34, 38, 39, 41]. Earlier return to daily activities and jogging may be explained by the effect of the transverse orientation of the femoral tunnel and graft using the AMP technique on the early postoperative period. We speculate that increased rotational stability in the early period after surgery enables aggressive rehabilitation and faster recovery.

This study has certain limitations. First, this was a retrospective study with no randomization. Second, it was a short-term study and thus could not assess the long-term results. Third, we did not assess knee stability objectively. Future randomized clinical trials directly comparing both techniques over long-term follow-up will help to clarify which technique provides the best clinical outcome.

In conclusion, the AMP technique is significantly superior to the TT technique in creating anatomical femoral tunnel placement during single-bundle ACL reconstruction. There is no evidence to support the superiority of either technique in terms of clinical outcomes. However, the AMP technique provides faster recovery in terms of return to normal life and return to jogging in the short-term.

## CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

## ACKNOWLEDGEMENTS

The authors thank Ibrahim Azboy, MD, for his contribution to this study.

## REFERENCES

- [1] Alentorn-Geli E, Samitier G, Alvarez P, Steinbacher G, Cugat R. Anteromedial portal versus transtibial drilling techniques in ACL reconstruction: a blinded cross-sectional study at two- to five-year follow-up. *Int Orthop* 2010; 34(5): 747-54.

- [http://dx.doi.org/10.1007/s00264-010-1000-1] [PMID: 20401753]
- [2] Saccomanni B. Graft fixation alternatives in anterior cruciate ligament reconstruction. *Musculoskelet Surg* 2011; 95(3): 183-91. [http://dx.doi.org/10.1007/s12306-011-0137-4] [PMID: 21538096]
- [3] Rayan F, Nanjayan SK, Quah C, Ramoutar D, Konan S, Haddad FS. Review of evolution of tunnel position in anterior cruciate ligament reconstruction. *World J Orthop* 2015; 6(2): 252-62. [http://dx.doi.org/10.5312/wjo.v6.i2.252] [PMID: 25793165]
- [4] Fu FH, van Eck CF, Tashman S, Irrgang JJ, Moreland MS. Anatomic anterior cruciate ligament reconstruction: a changing paradigm. *Knee Surg Sports Traumatol Arthrosc* 2015; 23(3): 640-8. [http://dx.doi.org/10.1007/s00167-014-3209-9] [PMID: 25086574]
- [5] Jepsen CF, Lundberg-Jensen AK, Faunoe P. Does the position of the femoral tunnel affect the laxity or clinical outcome of the anterior cruciate ligament-reconstructed knee? A clinical, prospective, randomized, double-blind study. *Arthroscopy* 2007; 23(12): 1326-33. [http://dx.doi.org/10.1016/j.arthro.2007.09.010] [PMID: 18063177]
- [6] Sadoghi P, Kröpfl A, Jansson V, Müller PE, Pietschmann MF, Fischmeister MF. Impact of tibial and femoral tunnel position on clinical results after anterior cruciate ligament reconstruction. *Arthroscopy* 2011; 27(3): 355-64. [http://dx.doi.org/10.1016/j.arthro.2010.08.015] [PMID: 21144694]
- [7] Carson EW, Anisko EM, Restrepo C, Panariello RA, O'Brien SJ, Warren RF. Revision anterior cruciate ligament reconstruction: etiology of failures and clinical results. *J Knee Surg* 2004; 17(3): 127-32. [PMID: 15366266]
- [8] Kamath GV, Redfern JC, Greis PE, Burks RT. Revision anterior cruciate ligament reconstruction. *Am J Sports Med* 2011; 39(1): 199-217. [http://dx.doi.org/10.1177/0363546510370929] [PMID: 20709943]
- [9] Andrei BI, Niculescu M, Popescu G. Position of anterior cruciate ligament after single-bundle arthroscopic reconstruction. *Int Orthop* 2016; 40(2): 393-7. [http://dx.doi.org/10.1007/s00264-015-2964-7] [PMID: 26330084]
- [10] Xu H, Zhang C, Zhang Q, *et al.* A systematic review of anterior cruciate ligament femoral footprint location evaluated by quadrant method for single-bundle and double-bundle anatomic reconstruction 2016; 32(8): 1724-34. [http://dx.doi.org/10.1016/j.arthro.2016.01.065]
- [11] Zhang Q, Zhang S, Li R, Liu Y, Cao X. Comparison of two methods of femoral tunnel preparation in single-bundle anterior cruciate ligament reconstruction: a prospective randomized study. *Acta Cir Bras* 2012; 27(8): 572-6. [http://dx.doi.org/10.1590/S0102-86502012000800010] [PMID: 22850710]
- [12] Zavras TD, Race A, Amis AA. The effect of femoral attachment location on anterior cruciate ligament reconstruction: graft tension patterns and restoration of normal anterior-posterior laxity patterns. *Knee Surg Sports Traumatol Arthrosc* 2005; 13(2): 92-100. [http://dx.doi.org/10.1007/s00167-004-0541-5] [PMID: 15756613]
- [13] Yagi M, Wong EK, Kanamori A, Debski RE, Fu FH, Woo SL. Biomechanical analysis of an anatomic anterior cruciate ligament reconstruction. *Am J Sports Med* 2002; 30(5): 660-6. [PMID: 12238998]
- [14] Duquin TR, Wind WM, Fineberg MS, Smolinski RJ, Buyea CM. Current trends in anterior cruciate ligament reconstruction. *J Knee Surg* 2009; 22(1): 7-12. [http://dx.doi.org/10.1055/s-0030-1247719] [PMID: 19216345]
- [15] Morgan CD, Kalman VR, Grawl DM. Definitive landmarks for reproducible tibial tunnel placement in anterior cruciate ligament reconstruction. *Arthroscopy* 1995; 11(3): 275-88. [http://dx.doi.org/10.1016/0749-8063(95)90003-9] [PMID: 7632302]
- [16] Reza zadeh S, Etehad H, Vosoughi AR. Outcome of arthroscopic single-bundle anterior cruciate ligament reconstruction: anteromedial portal technique *versus* transtibial drilling technique. *Musculoskelet Surg* 2016; 100(1): 37-41. [http://dx.doi.org/10.1007/s12306-015-0392-x] [PMID: 26645453]
- [17] Ferretti M, Ekdahl M, Shen W, Fu FH. Osseous landmarks of the femoral attachment of the anterior cruciate ligament: an anatomic study. *Arthroscopy* 2007; 23(11): 1218-25. [http://dx.doi.org/10.1016/j.arthro.2007.09.008] [PMID: 17986410]
- [18] Hefti F, Müller W, Jakob RP, Stäubli HU. Evaluation of knee ligament injuries with the IKDC form. *Knee Surg Sports Traumatol Arthrosc* 1993; 1(3-4): 226-34. [http://dx.doi.org/10.1007/BF01560215] [PMID: 8536037]
- [19] Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res* 1985; (198): 43-9. [PMID: 4028566]
- [20] Illingworth KD, Hensler D, Working ZM, Macalena JA, Tashman S, Fu FH. A simple evaluation of anterior cruciate ligament femoral tunnel position: the inclination angle and femoral tunnel angle. *Am J Sports Med* 2011; 39(12): 2611-8. [http://dx.doi.org/10.1177/0363546511420128] [PMID: 21908719]
- [21] Gadikota HR, Sim JA, Hosseini A, Gill TJ, Li G. The relationship between femoral tunnels created by the transtibial, anteromedial portal, and outside-in techniques and the anterior cruciate ligament footprint. *Am J Sports Med* 2012; 40(4): 882-8.

- [http://dx.doi.org/10.1177/0363546511434276] [PMID: 22302206]
- [22] Asagumo H, Kimura M, Kobayashi Y, Taki M, Takagishi K. Anatomic reconstruction of the anterior cruciate ligament using double-bundle hamstring tendons: surgical techniques, clinical outcomes, and complications. *Arthroscopy* 2007; 23(6): 602-9. [http://dx.doi.org/10.1016/j.arthro.2007.01.009] [PMID: 17560474]
- [23] Park JS, Park JH, Wang JH, et al. Comparison of femoral tunnel geometry, using *in vivo* 3-dimensional computed tomography, during transportal and outside-in single-bundle anterior cruciate ligament reconstruction techniques. *Arthroscopy* 2015; 31(1): 83-91. [http://dx.doi.org/10.1016/j.arthro.2014.07.023] [PMID: 25239169]
- [24] Dargel J, Schmidt-Wiethoff R, Fischer S, Mader K, Koebke J, Schneider T. Femoral bone tunnel placement using the transtibial tunnel or the anteromedial portal in ACL reconstruction: a radiographic evaluation. *Knee Surg Sports Traumatol Arthrosc* 2009; 17(3): 220-7. [http://dx.doi.org/10.1007/s00167-008-0639-2] [PMID: 18843479]
- [25] Silva A, Sampaio R, Pinto E. Placement of femoral tunnel between the AM and PL bundles using a transtibial technique in single-bundle ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2010; 18(9): 1245-51. [http://dx.doi.org/10.1007/s00167-010-1132-2] [PMID: 20390248]
- [26] Heming JF, Rand J, Steiner ME. Anatomical limitations of transtibial drilling in anterior cruciate ligament reconstruction. *Am J Sports Med* 2007; 35(10): 1708-15. [http://dx.doi.org/10.1177/0363546507304137] [PMID: 17664343]
- [27] Golish SR, Baumfeld JA, Schoderbek RJ, Miller MD. The effect of femoral tunnel starting position on tunnel length in anterior cruciate ligament reconstruction: a cadaveric study. *Arthroscopy* 2007; 23(11): 1187-92. [http://dx.doi.org/10.1016/j.arthro.2007.06.013] [PMID: 17986406]
- [28] Youm YS, Cho SD, Lee SH, Youn CH. Modified transtibial *versus* anteromedial portal technique in anatomic single-bundle anterior cruciate ligament reconstruction: comparison of femoral tunnel position and clinical results. *Am J Sports Med* 2014; 42(12): 2941-7. [http://dx.doi.org/10.1177/0363546514551922] [PMID: 25269655]
- [29] Azboy I, Demirtaş A, Gem M, Kiran S, Alemdar C, Bulut M. A comparison of the anteromedial and transtibial drilling technique in ACL reconstruction after a short-term follow-up. *Arch Orthop Trauma Surg* 2014; 134(7): 963-9. [http://dx.doi.org/10.1007/s00402-014-1996-6] [PMID: 24770982]
- [30] Noh JH, Roh YH, Yang BG, Yi SR, Lee SY. Femoral tunnel position on conventional magnetic resonance imaging after anterior cruciate ligament reconstruction in young men: transtibial technique *versus* anteromedial portal technique. *Arthroscopy* 2013; 29(5): 882-90. [http://dx.doi.org/10.1016/j.arthro.2013.01.025] [PMID: 23538044]
- [31] Sedeek SM, Andrew TH. Femoral tunnel drilling in anterior cruciate ligament reconstruction: anteromedial or transtibial portal? Current review. *Hard Tissue* 2013; 2: 34. [http://dx.doi.org/10.13172/2050-2303-2-4-718]
- [32] Riboh JC, Hasselblad V, Godin JA, Mather RC III. Transtibial *versus* independent drilling techniques for anterior cruciate ligament reconstruction: a systematic review, meta-analysis, and meta-regression. *Am J Sports Med* 2013; 41(11): 2693-702. [http://dx.doi.org/10.1177/0363546513506979] [PMID: 24130247]
- [33] Wei Z, Li F, Peng W, Wei B, Qiu L, Wei C. Comparative study on arthroscopic anterior cruciate ligament reconstruction with transtibial technique and through anteromedial approach. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 2014; 28(3): 339-44. [PMID: 24844016]
- [34] Mardani-Kivi M, Madadi F, Keyhani S, Karimi-Mobarake M, Hashemi-Motlagh K, Saheb-Ekhtiari K. Antero-medial portal *vs.* transtibial techniques for drilling femoral tunnel in ACL reconstruction using 4-strand hamstring tendon: a cross-sectional study with 1-year follow-up. *Med Sci Monit* 2012; 18(11): CR674-9. [http://dx.doi.org/10.12659/MSM.883546] [PMID: 23111744]
- [35] Robin BN, Jani SS, Marvil SC, Reid JB, Schillhammer CK, Lubowitz JH. Advantages and disadvantages of transtibial, anteromedial portal, and outside-in femoral tunnel drilling in single-bundle anterior cruciate ligament reconstruction: a systematic review. *Arthroscopy* 2015; 31(7): 1412-7. [http://dx.doi.org/10.1016/j.arthro.2015.01.018] [PMID: 25749530]
- [36] Musahl V, Voos JE, O'Loughlin PF, et al. Comparing stability of different single- and double-bundle anterior cruciate ligament reconstruction techniques: a cadaveric study using navigation. *Arthroscopy* 2010; 26(9)(Suppl.): S41-8. [http://dx.doi.org/10.1016/j.arthro.2010.01.028] [PMID: 20692120]
- [37] Marchant BG, Noyes FR, Barber-Westin SD, Fleckenstein C. Prevalence of nonanatomical graft placement in a series of failed anterior cruciate ligament reconstructions. *Am J Sports Med* 2010; 38(10): 1987-96. [http://dx.doi.org/10.1177/0363546510372797] [PMID: 20702859]
- [38] Bedi A, Musahl V, Steuber V, et al. Transtibial versus anteromedial portal reaming in anterior cruciate ligament reconstruction: an anatomic and biomechanical evaluation of surgical technique. *Arthroscopy* 2011; 27(3): 380-90. [http://dx.doi.org/10.1016/j.arthro.2010.07.018] [PMID: 21035990]
- [39] Bedi A, Raphael B, Maderazo A, Pavlov H, Williams RJ III. Transtibial *versus* anteromedial portal drilling for anterior cruciate ligament reconstruction: a cadaveric study of femoral tunnel length and obliquity. *Arthroscopy* 2010; 26(3): 342-50. [http://dx.doi.org/10.1016/j.arthro.2009.12.006] [PMID: 20206044]



- [40] Driscoll MD, Isabell GP Jr, Conditt MA, *et al.* Comparison of 2 femoral tunnel locations in anatomic single-bundle anterior cruciate ligament reconstruction: a biomechanical study. *Arthroscopy* 2012; 28(10): 1481-9. [<http://dx.doi.org/10.1016/j.arthro.2012.03.019>] [PMID: 22796141]
- [41] Tudisco C, Bisicchia S. Drilling the femoral tunnel during ACL reconstruction: transtibial *versus* anteromedial portal techniques. *Orthopedics* 2012; 35(8): e1166-72. [<http://dx.doi.org/10.3928/01477447-20120725-14>] [PMID: 22868600]

---

© Sukur *et al.*; Licensee Bentham Open

This is an open access article licensed under the terms of the Creative Commons Attribution-Non-Commercial 4.0 International Public License (CC BY-NC 4.0) (<https://creativecommons.org/licenses/by-nc/4.0/legalcode>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.