



The Open Orthopaedics Journal

Content list available at: <https://openorthopaedicsjournal.com>



REVIEW ARTICLE

Brace Prescription for Adult Scoliosis - Literature Review

Shu Yan Ng¹, Tai Hong Andrew Lung^{1*}, Lok Yan Joanne Cheng¹ and Yin Ling Elaine Ng²

¹Hong Kong Chiropractic College Foundation: 11&12/F China Hong Kong Tower, 8-12 Hennessy Road, Wan Chai, Hong Kong SAR

²Pedorthic Technology Limited, Wan Chai, Hong Kong SAR

Abstract:

Purpose:

The 2021 SOSORT guidelines stipulated that braces be prescribed for adult scoliosis with chronic pain and progressive curve. Yet, there have been no objective protocols relating to the prescription of the brace. Therefore, this review investigates if there are any objective criteria or generally agreed on protocols for brace prescription in adult scoliosis patients.

Methods:

Relevant papers were searched in PUBMED. Only articles that are in English and cover the clinical aspect of adult scoliosis bracing are included.

Results:

A total of twelve papers were identified. They include different adult scoliosis braces, ranging from elastic belts to rigid braces. The treatment protocol varied tremendously. No objective criteria were found concerning the prescription of a brace, daily wearing time, duration of the intervention, and weaning protocol. The brace treatment was primarily employed to manage low back pain.

Conclusion:

Our search showed no objective criteria and clear indications for bracing and no consensus concerning the prescription of braces, daily wearing time, and duration of the intervention for patients with adult scoliosis.

The authors proposed prescription of a brace be based on more objective radiological criteria and severity of low back pain. Brace prescription should depend on the flexibility of the curve and can range from accommodative to rigid braces of corrective design. "Corrective" brace has to be worn at least 14 hours daily for six months or until the low back pain subsides to the extent that permits daily activities with minimal discomfort. "Accommodative" brace can be worn when required.

Keywords: Adult idiopathic scoliosis, Degenerative lumbar scoliosis, Brace, Orthotic, Spine, Scoliosis.

Article History

Received: October 8, 2021

Revised: January 21, 2022

Accepted: March 23, 2022

1. INTRODUCTION

Adult scoliosis includes adult idiopathic scoliosis (IS) and degenerative lumbar scoliosis (DLS). Idiopathic scoliosis has its onset in childhood and adolescence, while degenerative lumbar scoliosis starts later in life, generally over 50. Thus, the prevalence of adult scoliosis is significantly higher than that of idiopathic scoliosis, particularly with increasing age, as the incidence of DLS increases with age [1, 2]. The reported prevalence of adult scoliosis ranges from 8.85% to 68% [3 - 5]. Fortunately, over 80% of the DLS patients have a mild curve of 10-20° [2].

Patients generally seek treatment for pain and disabilities rather than deformities [6]. Back pain is generally more common and severe in the thoracolumbar, lumbar, and lumbosacral curves than in the thoracic curve [7]. The thoracolumbar curve is treated surgically as the most common curve [8].

For symptomatic scoliosis patients, many different types of treatment have been studied. Unfortunately, dry needling, manipulation, and chiropractic treatment provide only short-term pain relief [9]. The temporary improvement is understandable, as the treatments address the pain instead of the underlying deranged spinal biomechanics. Adult scoliosis brace, which attempts to stabilize or improve the altered spinal biomechanics, has a better outcome but minimal evidence [9].

* Address correspondence to this author at the Hong Kong Chiropractic College Foundation, 11&12/F China Hong Kong Tower, 8-12 Hennessy Road, Wan Chai, Hong Kong SAR; Tel: +852 39983208; Fax: +852 39983222; E-mail: lthandy@outlook.com

A recent systematic literature review concluded that many bracing studies on adult scoliosis patients are biased [10].

However, recent studies have shown that bracing provides pain relief and stabilizes postures [11 - 13]. Also, the 2011 SOSORT guidelines stipulated that braces be prescribed for adult scoliosis patients with chronic pain and progressive scoliosis curves [14].

Given the inconclusive evidence, clinicians sometimes have difficulty determining if a brace is indicated in adult scoliosis patients presented with low back pain. We thus undertake to review the literature to determine if there are any objective criteria upon which we can base when prescribing an adult scoliosis brace. We also investigate whether there is a generally agreed protocol for brace wear in adult scoliosis patients.

2. MATERIALS AND METHODS

Relevant papers were searched in PUBMED, using the Boolean search operators: (“adult scoliosis” OR “adult spinal deformity”) AND (“orthotic” OR “brace”), and the keywords “adult scoliosis” AND “radiological parameters.” All titles were then screened for relevancy. Papers not in English and irrelevant to the present investigation were excluded. Only those papers, including brace as treatment and studying the radiological parameters of adult scoliosis, were reviewed. Also, a manual search was undertaken from the references of the selected papers.

3. RESULTS

The search yielded 17 papers, of which 12 were regarded as appropriate. In addition, a manual search of the articles revealed other articles on scoliosis curve progression.

A review of the bracing studies showed that most braces were prescribed for adult scoliosis patients with thoracolumbar, lumbar, and double curves. The Cobb angle ranged from 10° to 91° [11 - 13], with a visual analog scale (VAS) of low back pain of not less than 6/10 [11, 15, 16]. Only one study included patients with the thoracic curve [17]. Most of the subjects had chronic low back pain.

Different types of braces have been used for the treatment. They ranged from the modified off-the-shelf elastic belt [16], a specific scoliosis soft brace (Peak Scoliosis Brace) [17, 18], Spinecor [19], to rigid braces [11 - 13, 20, 21]. Of interest is that the rigid braces prescribed employed different biomechanical principles. The Physiologic[®] brace is a sagittal alignment brace used to restore sagittal alignment in adult scoliosis patients. The sBrace [15], Vesinet TLSO [13], the Lyon brace [12, 21], and the ART brace [12] attempted to correct or accommodate scoliosis in three dimensions. However, no standards have been found concerning the prescription of the braces. The Peak Scoliosis brace, which is an off-the-shelf brace, was prescribed for adult scoliosis patients, with a mean Cobb angle of 61.9° [17], while rigid braces were prescribed for curves, with Cobb angle measuring 36—50° [11 - 13, 21] (Table 1).

Table 1. The intervention and outcome measures of different brace studies, which used soft braces and rigid braces.

Scoliosis	Paper	Study	Age	No/Sex	Complaints		Brace	Prescription	Outcome		Remarks
					LBP: VAS	Curve			VAS	Cobb	
Adult AIS	Marcotte 2010 [19]	Prosp	18-69	26F; 4M	-	-	Spinecor	10-130 hrs/wk; 18-28 mths	Improved 77%	-	-
	Weiss <i>et al.</i> 2016 [20]	Case Report	37	1F	Average: 5-7/10 Occasional episodes: 8-9/10	Double curve; Th 56°; LB 50°	Gensingen	3-4 hrs daily; 3 days/wk 16 mths	0/10	Th/L: 55/32	Ex
	Polastri and Romano 2017 [16]	Case Report	40	1F	8.5/10	ThL and LB curves: 22°	Customized elastic brace	At-will; 24 mths	2/10	-	-
	de Mauroy <i>et al.</i> 2016 [12]	Prosp	56 ± 17	144F; 14M	-	ThL and LB: 39.7 ± 17.4°	Plaster cast + Lyon brace	3 wks plaster; 6 mths Lyon brace 4 hrs/day; ≥ 5 yrs		24% improved > 5°; 56% stable; 20% worsened > 5°	Ex; Cobb angle: NS Coronal Bal: S Sagittal Bal: NS
	Han <i>et al.</i> , 2020 [23]	Prosp	24.3	17F; 1M	-	29.3°	All Line	12 hrs/day; 12 wks	-	24.0°	Sig reduction in Cobb angle
DLS	Weiss and Dallmayer 2006 [22]	Case Report	47	1F	8000 steps medicated	LB: 55°; LB kyphosis: 30°	Physiologic Brace	Daily: at-will 10 days	Reduce by 1/5; 12000 steps	-	-

Adult AIS and DLS	Weiss <i>et al</i> 2006 [24]	Prosp	41±21	29F	3.38/5	37 ± 22°	Physiologic Brace	>8 hrs/day (at will); av 7.5 mths	2.69/5	-	Poor compliance
	Weiss and Werkmann 2009 [11]	Prosp	NA	58F; 9M	3.3/5	41 ± 22° (10-91 °)	Physiologic Brace	20hr/day x 6 mths; case by case; 18 mths	2.0/5	-	Changes in pain measures: Sig
	Gallo 2014 [15]	Case Reports	C1: 65 C2: NA	2F	C1: 8-9/10; C2: 8/10	NA	sBrace	At-will	C1: 3-4/10; C2: 2/10	-	-
	de Mauroy <i>et al</i> 2016 [21]	Prosp	57±16	650F; 89M	-	36.0 ± 17.4°	Plaster cast + Lyon brace	3 wks plaster; 6 mths Lyon brace; 15 yrs	-	-	Non-adherent: 17%
	Palazzo <i>et al</i> 2017 [13]	Retrosp	61±8	38F	-	49.6±17.7°; Progression rate: 1.28/yr	Vesinet TLSO	≥ 6 hrs/day; 5 yrs	Progression rate: 0.21/yr	-	-
	Zaina <i>et al</i> 2018 [17]	Prosp	68±11	20F	7.15/10	Av Cobb: 61.9° 55% ThL/LB; 30% Double curve; 15% Th	Peak Scoliosis Brace	2-4 hrs/day; 4 wks	5.85/10	-	-

Th stands for thoracic; L and LB stand for lumbar; S and sig. stand for significant; NS stands for non-significant; Prosp stands for prospective; Retrosp stands for retrospective; Ex stands for exercise; NA stands for not available; C1 stands for Case One; C2 stands for Case two.

Also, the daily wear of the brace varied tremendously with the studies, ranging from at-will [15, 22] to 12 [23] to 20 hours/day [11], and 24 hours/day [12]. The variation may be related to the severity of the structural curves and the intensity of pain. De Mauroy *et al.* [21] reasoned that the extended period of daily wear was necessary to elongate the contracted tissues.

Further, the duration of the intervention is not uniform. Palazzo *et al.* [13] and de Mauroy *et al.* [21] did not describe the length of treatment; they followed up the patients for more than 5 and 15 years, respectively. Weiss and Werkmann [11] reported assessing the patient again 18 months after the initial brace wear.

There were no studies on in-brace correction in adult scoliosis. The outcome measured was on reduction of progression rate [12, 13] and pain intensity, which are the most common and easiest measures of brace success [11, 15, 17]. De Mauroy *et al.* [12] showed that the Lyon brace improved 24% of the curves by ≥5°, stabilized 56% of the curves, and worsened 20% by ≥5° in 158 patients with adult scoliosis. Similarly, Palazzo *et al.* [13] reported that brace treatment reduced the progression of adult scoliosis curves from 1.28° to 0.21° per year.

4. DISCUSSION

The studies have shown that rigid and soft braces reduce low back pain, and rigid brace contains curve progression in adult scoliosis patients. The indications are consistent with the SOSORT guidelines, which stipulated the brace prescription when back pain and curve progression are maximal in adult scoliosis patients [14]. A brace is indicated for patients whose low back pain did not respond to conventional treatments [11, 12, 21]. It is also used to stabilize adult scoliosis [12, 21] and reduce the curve progression rate [13]. In a retrospective cohort study that involved 38 female patients, a spinal brace reduces the curve progression rate from 1.28° to 0.21 ° per year [13].

The daily brace wearing hours vary with studies, from “at-will” [15] to 24-hours [12, 21]. In addition, the wear duration ranges from 7.5 months [24] to 15 years [21]. The studies, however, did not describe the weaning protocols.

Overall, the studies did not specify any objective criteria for brace prescription for adult scoliosis patients. Therefore, there are no generally agreed-on prescription protocols for braces for adult scoliosis patients. We thus attempt to propose a protocol for adult scoliosis bracing based on more objective radiological parameters.

Many radiological parameters have been found to associate with low back pain. Low back pain is more frequently associated with double scoliosis, thoracolumbar and lumbar curves, and less often with the thoracic curve. Lumbar rotatoryolisthesis over 5mm [25, 26], thoracolumbar or lumbar curve over 30°, apical lumbar vertebral rotation over 33%, and a low intercrystal line [27] are associated with more frequent low back pain and radiculopathy. Recently, Ferrero *et al.* [28] showed that DLS patients with axial intervertebral rotation (AIR) at apex over 10° had significantly worse Oswestry Disability index (ODI) and more low back pain. The association is understandable as the AIR in the upper and lower junctions of the scoliosis curve subject the curve to torque (Fig. 1), increasing lateral scoliotic deviation [28].

Sagittal radiological parameters have also been found to correlate with disability. A pelvic tilt >22° and pelvic incidence-lumbar lordosis >10° are associated with disability [29], suggesting pelvic retroversion. The sagittal vertical axis (SVA), which is the distance between a plumb line dropped from C7 to the sacrum's posterosuperior angle, is also associated with clinical symptoms; the threshold relating to clinical symptoms is variably reported to be 43mm [29] and 70mm [30]. The increased SVA indicates a truncal forward shift relative to the pelvis to increase the size of the spinal canal and reduce neurologic insult, especially in the presence of degenerative spondylolisthesis [31]. Yet, it is of note that

SVA can be diminished by pelvic retroversion. Thus, SVA and pelvic retroversion must be considered together [32]. T1 pelvic angle, which integrates the forward trunk inclination and the pelvis compensation, may be a better parameter than SVA [33]. It is less influenced by posture [33] and has been found to correlate strongly with the SVA, pelvic tilt, and pelvic incidence lumbar lordosis mismatch (PI-LL) [34].




Fig. (1). The measurement of axial intervertebral rotation. It is the relative intervertebral rotation between adjacent vertebrae within the scoliosis curve.

Some of the above radiological parameters may correlate with curve progression. A lumbar rotatoryolisthesis of more than 6mm, thoracolumbar or lumbar curve $>30^\circ$, apical vertebral rotation $>33\%$, and a low intercrystal line signify a higher propensity for scoliosis curve progression [35 - 38]. A recent study showed that the magnitude of the Cobb angle and the level of the intercrystal line bear no relationship to the risk of curve progression in patients with adult spinal deformity [39]. However, a truncal shift of 5mm to the left of the central sacral line (CSL) in the presence of a left thoracolumbar or lumbar curve is a risk factor for curve progression [40]. The curve progression is associated with a significant coronal shift of L1 from CSL [41].

In many instances, patients do not have prior radiographs.

Documentation of the rate of progression of the curve as a justification for bracing would thus be challenging, especially when adult IS patients have a long history of scoliosis, and progression over the years may not be linear [42, 43]. Therefore, it is proposed that adult scoliosis patients with sagittal imbalance, reduced lumbar lordosis, and radiological parameters associated with a higher risk of curve progression or low back pain are indicated for brace treatment (Fig. 2). It has to be noted that low back pain in a scoliotic patient does not equate that the pain must be related to scoliosis [44]. Before the brace prescription, a proper evaluation should ensure that the pain is secondary to the spinal deformity and not other causes.

Whether the brace should be accommodative or corrective depends on the deformities and spinal rigidity and not on the patients' body habitus. At present, it is clinically challenging to grade lumbar rigidity. The presence of marked arthritic or osteochondral changes suggests an increased spinal rigidity. An accommodative brace should be prescribed when the curve is rigid [17, 18] or in the presence of rotatoryolisthesis to help stabilize the spine (Fig. 3). When the elastic brace cannot provide the needed support because of a mismatch of the spinal contour, a custom accommodative rigid brace may be prescribed to provide spinal support and limit motion, particularly in patients with more severe curves. Coupled movements in the lumbar spines are higher in DLS patients, particularly with larger curves, in most loading directions than in the lumbar spine with no scoliosis [45].

When the scoliotic deformity is rigid coronally but flexible in the sagittal plane, a rigid two-dimensional sagittal realignment brace may be prescribed to improve sagittal balance while accommodating deformities in the coronal and transverse planes. The sagittal spinal parameters worsen in patients with ASD with age; the change predominantly involves reducing lumbar lordosis, though the reduction is not statistically significant [39].

However, when the scoliosis curve is flexible, as in young adults and peri-menopausal patients, a 3D rigid brace with a corrective design may be prescribed.

The duration of daily wear depends on the severity of the symptoms and the extent of the spinal deformities. Weiss *et al.* [24] showed that brace wear 4-8 hours daily did not significantly improve low back pain. However, an increase in dose to 20 hours daily significantly improved the outcome [11]. De Mauroy *et al.* [12, 21] reported similar findings, who braced the patients for 24 hours per day in the first three weeks using a plaster cast and more than 4 hours daily after that. In the presence of radiological parameters indicative of curve progression, loss of lumbar lordosis, sagittal spinal imbalance, and increased risk of low back pain, the authors suggest wearing the brace during the daytime, for at least 14 hours a day, to reduce the axial deforming loading of body weight on the spine [46]. It has to be noted that the hours of wear were reported by patients and were not measured objectively by thermal or pressure sensors.




Fig. (2). The proposed brace type prescription for adult scoliosis patients. PSSE refers to physiotherapeutic scoliosis specific exercise; Th/L refers to thoracolumbar region; PI refers to pelvic incidence; LL refers to lumbar lordosis; PT refers to pelvic tilt.




Fig. (3). (a) An adult with idiopathic scoliosis. She is in her thirties, and rigid brace if prescribed can be corrective in three dimensional. (b) An adult with degenerative lumbar scoliosis. Rotary olisthesis is evident at L3/4 level, with coronal shift. (c) The lateral full spine x-ray of the same patient as (b). Prescription of a rigid brace in this patient should be a sagittal realignment brace.

As in brace treatment of adolescent IS [47], daily core strengthening exercises should be prescribed to avoid deconditioning the lumbar paraspinal muscles and postural collapse during brace wear and after brace weaning, as sarcopenia is a risk factor in the progression of adult scoliosis. In addition, patients should learn diaphragmatic and de-rotational breathing to strengthen the deep lumbar muscles, enhance spinal stability, and reduce vertebral rotation and torque from the scoliosis curve [48, 49]. Also, they have to perform exercises to improve the spine's sagittal alignment and adopt proper sitting postures to reduce lumbar intervertebral disc pressure [50]. Treatment should also be given for osteopenia or osteoporosis [51].

Weiss *et al.* [24] and de Mauroy *et al.* [12, 21] reported having patients wearing the brace for six months before weaning. Our clinical experience concurred with the findings; we propose that the corrective brace be worn for 14 hours per day for at least six months or until the low back pain reduces to a level the patient can engage in daily activity with minimal pain before weaning. The wear time can be reduced by two hours every two months until the patient wears the brace for only 8 hours a day. A lateral spinal radiograph is then taken to determine the sagittal parameters. When no improvement is evident, the patient should not further reduce the daily brace wear; further weaning should depend on the outcome of the clinical evaluation. In the presence of sagittal parameters improvement, the brace wear can be further reduced by two hours every two months until the brace is weaned off completely. The authors are not aware of any studies showing changes in sagittal alignment with braces. However, the importance of improving the sagittal spinal parameters was brought forward by Pizones *et al.* [52]. A study on adult scoliosis surgery showed that the surgery should restore the ideal Rousouly sagittal profile associated with decreased mechanical complications, especially in patients over 65 [52]. In addition, the brace should target restoring the spinopelvic relationship and zero out the compensatory mechanism, improving the lumbar lordosis and pelvic tilt [53]. On the other hand, the accommodative brace is proposed to be worn when required.

The proposed clinical guidelines for adult scoliosis brace have limitations [10]. They are empirical and are not evidence-based. More studies are required to establish the effectiveness and dosage of adult bracing to manage chronic low back pain in adult scoliosis patients. Subjects should preferably be more homogenous, as the response of adult IS patients to bracing may be very much different from that of DLS patients, and different curve types respond to bracing differently.

CONCLUSION

Our search showed no objective criteria or clear indications concerning the prescription of the types of braces, daily wearing time, and duration of the intervention for adult scoliosis. Therefore research is required to determine the effectiveness and protocols of bracing in managing adult scoliosis patients, particularly those with thoracolumbar curves.

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Kilshaw M, Baker RP, Gardner R, Charosky S, Harding I. Abnormalities of the lumbar spine in the coronal plane on plain abdominal radiographs. *Eur Spine J* 2011; 20(3): 429-33. [http://dx.doi.org/10.1007/s00586-010-1610-8] [PMID: 21069544]
- [2] Xu L, Sun X, Huang S, *et al.* Degenerative lumbar scoliosis in Chinese Han population: prevalence and relationship to age, gender, bone mineral density, and body mass index. *Eur Spine J* 2013; 22(6): 1326-31. [http://dx.doi.org/10.1007/s00586-013-2678-8] [PMID: 23361532]
- [3] Kebaish KM, Neubauer PR, Voros GD, Khoshnevisan MA, Skolasky RL. Scoliosis in adults aged forty years and older: prevalence and relationship to age, race, and gender. *Spine* 2011; 36(9): 731-6. [http://dx.doi.org/10.1097/BRS.0b013e3181e9f120] [PMID: 20881515]
- [4] Kobayashi T, Atsuta Y, Takemitsu M, Matsuno T, Takeda N. A prospective study of *de novo* scoliosis in a community based cohort. *Spine* 2006; 31(2): 178-82. [http://dx.doi.org/10.1097/01.brs.0000194777.87055.1b] [PMID: 16418637]
- [5] Schwab F, Dubey A, Pagala M, Gamez L, Farcy JP. Adult scoliosis: a health assessment analysis by SF-36. *Spine* 2003; 28(6): 602-6. [http://dx.doi.org/10.1097/01.BRS.0000049924.94414.BB] [PMID: 12642769]
- [6] Nakamae T, Yamada K, Shimbo T, *et al.* Bone marrow edema and low back pain in elderly degenerative lumbar scoliosis. *Spine* 2016; 41(10): 885-92. [http://dx.doi.org/10.1097/BRS.0000000000001315] [PMID: 26641841]
- [7] Jackson RP, Simmons EH, Stripinis D. Incidence and severity of back pain in adult idiopathic scoliosis. *Spine* 1983; 8(7): 749-56. [http://dx.doi.org/10.1097/00007632-198310000-00011] [PMID: 6229884]
- [8] Pizones J, Pérez Martín-Buitrago M, Perez-Grueso FJS, *et al.* Function and clinical symptoms are the main factors that motivate thoracolumbar adult scoliosis patients to pursue surgery. *Spine* 2017; 42(1): E31-6. [http://dx.doi.org/10.1097/BRS.0000000000001694] [PMID: 27196023]
- [9] Everett CR, Patel RK. A systematic literature review of nonsurgical treatment in adult scoliosis. *Spine* 2007; 32(19)(Suppl.): S130-4. [http://dx.doi.org/10.1097/BRS.0b013e318134ea88] [PMID: 17728680]
- [10] McAviney J, Mee J, Fazalbhoy A, Du Plessis J, Brown BT. A systematic literature review of spinal brace/orthosis treatment for adults with scoliosis between 1967 and 2018: clinical outcomes and harms data. *BMC Musculoskelet Disord* 2020; 21(1): 87. [http://dx.doi.org/10.1186/s12891-020-3095-x] [PMID: 32035480]
- [11] Weiss HR, Werkmann M. Treatment of chronic low back pain in patients with spinal deformities using a sagittal re-alignment brace. *Scoliosis* 2009; 4(1): 7. [http://dx.doi.org/10.1186/1748-7161-4-7] [PMID: 19272146]
- [12] de Mauroy JC, Lecante C, Barral F, Pourret S. Prospective study of 158 adult scoliosis treated by a bivalve polyethylene overlapping brace and reviewed at least 5 years after brace fitting. *Scoliosis Spinal Disord* 2016; 11(S2)(Suppl. 2): 28. [http://dx.doi.org/10.1186/s13013-016-0091-x] [PMID: 27785467]
- [13] Palazzo C, Montigny JP, Barbot F, *et al.* Effects of bracing in adult

- with scoliosis: a retrospective study. *Arch Phys Med Rehabil* 2017; 98(1): 187-90.
[<http://dx.doi.org/10.1016/j.apmr.2016.05.019>] [PMID: 27343345]
- [14] Negrini S, Donzelli S, Aulisa AG, *et al.* 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis Spinal Disord* 2018; 13(1): 3.
[<http://dx.doi.org/10.1186/s13013-017-0145-8>] [PMID: 29435499]
- [15] Gallo D. Case reports: orthotic treatment of adult scoliosis patients with chronic back pain. *Scoliosis* 2014; 9(1): 18.
[<http://dx.doi.org/10.1186/1748-7161-9-18>] [PMID: 25904972]
- [16] Polastri M, Romano M. Lumbar scoliosis: Reducing lower back pain and improving function in adulthood. A case report with a 2-year follow-up. *J Bodyw Mov Ther* 2017; 21(1): 81-5.
[<http://dx.doi.org/10.1016/j.jbmt.2016.05.004>] [PMID: 28167195]
- [17] Zaina F, Poggio M, Donzelli S, Negrini S. Can bracing help adults with chronic back pain and scoliosis? Short-term results from a pilot study. *Prosthet Orthot Int* 2018; 42(4): 410-4.
[<http://dx.doi.org/10.1177/0309364618757769>] [PMID: 29446692]
- [18] Zaina F, Poggio M, Di Felice F, Donzelli S, Negrini S. Bracing adults with chronic low back pain secondary to severe scoliosis: six months results of a prospective pilot study. *Eur Spine J* 2021; 30(10): 2962-6.
[<http://dx.doi.org/10.1007/s00586-021-06808-1>] [PMID: 33733328]
- [19] Marcotte L. SpineCor in the treatment of adult scoliosis. *Scoliosis* 2010; 5(S1): O47.
[<http://dx.doi.org/10.1186/1748-7161-5-S1-O47>]
- [20] Weiss HR, Moramarco K, Moramarco M. Scoliosis bracing and exercise for pain management in adults-a case report. *J Phys Ther Sci* 2016; 28(8): 2404-7.
[<http://dx.doi.org/10.1589/jpts.28.2404>] [PMID: 27630444]
- [21] de Mauroy JC, Lecante C, Barral F, Pourret S. Bracing in adult with scoliosis: experience in diagnosis and classification from a 15 year prospective study of 739 patients. *Scoliosis Spinal Disord* 2016; 11(S2)(Suppl. 2): 29.
[<http://dx.doi.org/10.1186/s13013-016-0090-y>] [PMID: 27785468]
- [22] Weiss HR, Dallmayer R. Brace treatment of spinal claudication in an adult with lumbar scoliosis--a case report. *Stud Health Technol Inform* 2006; 123: 586-9.
[PMID: 17108492]
- [23] Han KS, Kim GW, Kang SR, Ko MH, Seo JH. Clinical evaluation of the effectiveness of a new orthotic device for the non-operative treatment of scoliosis. *Technol Health Care* 2020; 28(S1): 229-36.
[<http://dx.doi.org/10.3233/THC-209023>] [PMID: 32364155]
- [24] Weiss HR, Dallmayer R, Stephan C. First results of pain treatment in scoliosis patients using a sagittal realignment brace. *Stud Health Technol Inform* 2006; 123: 582-5.
[PMID: 17108491]
- [25] Marty-Poumarat C, Scattin L, Marpeau M, Garreau de Loubresse C, Aegerter P. Natural history of progressive adult scoliosis. *Spine* 2007; 32(11): 1227-34.
[<http://dx.doi.org/10.1097/01.brs.0000263328.89135.a6>] [PMID: 17495780]
- [26] Ferrero E, Lafage R, Chailly V, *et al.* Clinical and stereoradiographic analysis of adult spinal deformity with and without rotatory subluxation. *Orthop Traumatol Surg Res* 2015; 101(5): 613-8.
[<http://dx.doi.org/10.1016/j.otsr.2015.04.008>] [PMID: 26194209]
- [27] Weinstein SL, Ponseti IV. Curve progression in idiopathic scoliosis. *J Bone Joint Surg Am* 1983; 65(4): 447-55.
[<http://dx.doi.org/10.2106/00004623-198365040-00004>] [PMID: 6833318]
- [28] Ferrero E, Lafage R, Diebo BG, *et al.* Tridimensional analysis of rotatory subluxation and sagittal spinopelvic alignment in the setting of adult spinal deformity. *Spine Deform* 2017; 5(4): 255-64.
[<http://dx.doi.org/10.1016/j.jspd.2017.01.003>] [PMID: 28622901]
- [29] Schwab FJ, Blondel B, Bess S, *et al.* Radiographical spinopelvic parameters and disability in the setting of adult spinal deformity: a prospective multicenter analysis. *Spine* 2013; 38(13): E803-12.
[<http://dx.doi.org/10.1097/BRS.0b013e318292b7b9>] [PMID: 23722572]
- [30] Glassman SD, Berven S, Bridwell K, Horton W, Dimar JR. Correlation of radiographic parameters and clinical symptoms in adult scoliosis. *Spine* 2005; 30(6): 682-8.
[<http://dx.doi.org/10.1097/01.brs.0000155425.04536.f7>] [PMID: 15770185]
- [31] Buckland AJ, Vira S, Oren JH, *et al.* When is compensation for lumbar spinal stenosis a clinical sagittal plane deformity? *Spine J* 2016; 16(8): 971-81.
[<http://dx.doi.org/10.1016/j.spinee.2016.03.047>] [PMID: 27063925]
- [32] Lafage V, Schwab F, Patel A, Hawkinson N, Farcy JP. Pelvic tilt and truncal inclination: two key radiographic parameters in the setting of adults with spinal deformity. *Spine* 2009; 34(17): E599-606.
[<http://dx.doi.org/10.1097/BRS.0b013e3181aad219>] [PMID: 19644319]
- [33] Protosaltis T, Schwab F, Bronsard N, *et al.* The T1 pelvic angle, a novel radiographic measure of global sagittal deformity, accounts for both spinal inclination and pelvic tilt and correlates with health-related quality of life. *J Bone Joint Surg Am* 2014; 96(19): 1631-40.
[<http://dx.doi.org/10.2106/JBJS.M.01459>] [PMID: 25274788]
- [34] Banno T, Hasegawa T, Yamato Y, *et al.* T1 pelvic angle is a useful parameter for postoperative evaluation in adult spinal deformity patients. *Spine* 2016; 41(21): 1641-8.
[<http://dx.doi.org/10.1097/BRS.0000000000001608>] [PMID: 27802254]
- [35] Kelly MP, Lurie JD, Yanik EL, *et al.* Operative versus nonoperative treatment for adult symptomatic lumbar scoliosis. *J Bone Joint Surg Am* 2019; 101(4): 338-52.
[<http://dx.doi.org/10.2106/JBJS.18.00483>] [PMID: 30801373]
- [36] Seo JY, Ha KY, Hwang TH, Kim KW, Kim YH. Risk of progression of degenerative lumbar scoliosis. *J Neurosurg Spine* 2011; 15(5): 558-66.
[<http://dx.doi.org/10.3171/2011.6.SPINE10929>] [PMID: 21780861]
- [37] Silva FE, Lenke LG. Adult degenerative scoliosis: evaluation and management. *Neurosurg Focus* 2010; 28(3): E1.
[<http://dx.doi.org/10.3171/2010.1.FOCUS09271>] [PMID: 20192655]
- [38] Pritchett JW, Bortel DT. Degenerative symptomatic lumbar scoliosis. *Spine* 1993; 18(6): 700-3.
[<http://dx.doi.org/10.1097/00007632-199305000-00004>] [PMID: 8516697]
- [39] Faraj SSA, Te Hennepe N, van Hooff ML, Pouw M, de Kleuver M, Spruit M. The natural history of progression in adult spinal deformity: a radiographic analysis. *Global Spine J* 2020; 10(3): 272-9.
[<http://dx.doi.org/10.1177/2192568219845659>] [PMID: 32313792]
- [40] Ushirozako H, Yoshida G, Hasegawa T, *et al.* Impact of shift to the concave side of the C7-center sacral vertical line on de novo degenerative lumbar scoliosis progression in elderly volunteers. *J Orthop Sci* 2020; 25(1): 82-8.
[<http://dx.doi.org/10.1016/j.jos.2019.03.007>] [PMID: 30926295]
- [41] Tsutsui S, Yoshimura N, Watanuki A, *et al.* Risk factors and natural history of de novo degenerative lumbar scoliosis in a community-based cohort: the Miyama study. *Spine Deform* 2013; 1(4): 287-92.
[<http://dx.doi.org/10.1016/j.jspd.2013.05.005>] [PMID: 27927360]
- [42] Asceni E, Bartolozzi P, Logroscino CA, *et al.* Natural history of untreated idiopathic scoliosis after skeletal maturity. *Spine* 1986; 11(8): 784-9.
[<http://dx.doi.org/10.1097/00007632-198610000-00007>] [PMID: 3810293]
- [43] Kotwal S, Pumberger M, Hughes A, Girardi F. Degenerative scoliosis: a review. *HSS J* 2011; 7(3): 257-64.
[<http://dx.doi.org/10.1007/s11420-011-9204-5>] [PMID: 23024623]
- [44] Janicki JA, Alman B. Scoliosis: Review of diagnosis and treatment. *Paediatr Child Health* 2007; 12(9): 771-6.
[<http://dx.doi.org/10.1093/pch/12.9.771>] [PMID: 19030463]
- [45] Rustenburg CME, Kingma I, Holewijn RM, *et al.* Biomechanical properties in motion of lumbar spines with degenerative scoliosis. *J Biomech* 2020; 102: 109495.
[<http://dx.doi.org/10.1016/j.jbiomech.2019.109495>] [PMID: 31767285]
- [46] Fei H, Li WS, Sun ZR, Jiang S, Chen ZQ. Effect of patient position on the lordosis and scoliosis of patients with degenerative lumbar scoliosis. *Medicine (Baltimore)* 2017; 96(32): e7648.
[<http://dx.doi.org/10.1097/MD.0000000000007648>] [PMID: 28796046]
- [47] Negrini S, Zaina F, Romano M, Negrini A, Parzini S. Specific exercises reduce brace prescription in adolescent idiopathic scoliosis: a prospective controlled cohort study with worst-case analysis. *J Rehabil Med* 2008; 40(6): 451-5.
[<http://dx.doi.org/10.2340/16501977-0195>] [PMID: 18509560]
- [48] Kolar P, Sulc J, Kyncl M, *et al.* Postural function of the diaphragm in persons with and without chronic low back pain. *J Orthop Sports Phys Ther* 2012; 42(4): 352-62.
[<http://dx.doi.org/10.2519/jospt.2012.3830>] [PMID: 22236541]
- [49] Lehnert-Schroth C, Grobl P. *Dreidimensionale skoliosebehandlung*. 8th ed. Munich: Elsevier GmbH 2014. Three-dimensionally treatment for scoliosis
- [50] Wilke HJ, Neef P, Caimi M, Hoogland T, Claes LE. New *in vivo*

- measurements of pressures in the intervertebral disc in daily life. *Spine* 1999; 24(8): 755-62.
[<http://dx.doi.org/10.1097/00007632-199904150-00005>] [PMID: 10222525]
- [51] Ng SY. Conservative treatment of degenerative lumbar scoliosis. *Spinal Deformities in Adolescents, Adults and Older Adults*. IntechOpen 2019.
- [52] Pizones J, Moreno-Manzanaro L, Sánchez Pérez-Grueso FJ, *et al.* Restoring the ideal Roussouly sagittal profile in adult scoliosis surgery decreases the risk of mechanical complications. *Eur Spine J* 2020; 29(1): 54-62.
[<http://dx.doi.org/10.1007/s00586-019-06176-x>] [PMID: 31641904]
- [53] Savarese LG, Menezes-Reis R, Bonugli GP, Herrero CFPDS, Defino HLA, Nogueira-Barbosa MH. Spinopelvic sagittal balance: what does the radiologist need to know? *Radiol Bras* 2020; 53(3): 175-84.
[<http://dx.doi.org/10.1590/0100-3984.2019.0048>] [PMID: 32587427]

© 2022 Ng *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.