

Reverse Shoulder Arthroplasty in Rheumatoid Arthritis: A Systematic Review

Edward C.A. Gee*, Emma K. Hanson and Adnan Saithna

Southport and Ormskirk Hospital NHS Trust, Town Lane, Southport, Merseyside, PR8 6PN, UK

Abstract: *Background:* Anatomical shoulder replacement for rheumatoid arthritis (RA) is complicated by a high incidence of rotator cuff tears and glenoid erosion. This can lead to poor function and early failure. Reverse shoulder arthroplasty (RSA) has gained popularity as an alternative. This systematic review attempts to further define the role of RSA in RA.

Methods: A systematic review identified seven studies reporting outcomes of RSA in RA patients. Studies were critically appraised, and data on outcomes, complications and technical considerations were extracted and analysed.

Results: One hundred and twenty one shoulders were included (mean follow up 46.9 months). Consistent improvements in the main outcome measures were noted between studies. Ninety five percent of patients described excellent to satisfactory outcomes. The minimum mean forward elevation reported in each study was 115 degrees. Symptomatic glenoid loosening (1.7%), deep infection (3.3%) and revision surgery (5%) rates were no higher than for a population of mixed aetiologies.

Discussion: Previous concerns regarding high pre- and peri-operative complication and revision rates in RA patients were not shown to be valid by the results of this review. Although associated cuff tears are common and glenoid bone loss can increase the technical complexity of surgery, RSA provides consistent and predictable improvements in key outcome measures and the revision and complication rates do not appear to be higher than reported in a large population of mixed aetiologies.

Conclusion: The contemporary literature shows that RSA is a safe, effective and reliable treatment option in RA patients.

Keywords: Outcomes, reverse shoulder arthroplasty, rheumatoid arthritis.

INTRODUCTION

The glenohumeral joint is frequently affected in rheumatoid arthritis (RA) but the incidence is variably reported. Cuomo *et al.* found that 90% of patients with RA had some shoulder impairment. However, this report predates the significant improvements seen in the medical management of RA in the last decade [1]. More recently, Lehtinen *et al.* reported that 48% of RA patients developed glenohumeral erosive changes and Rittmeister *et al.* reported that 75% of those with seropositive RA developed shoulder symptoms [2, 3]. Despite this variability, there is no doubt that shoulder involvement is common in RA. In those patients who fail non-operative management, arthroplasty surgery is an option. However, there are several additional concerns when RA patients require shoulder replacement compared to those with glenohumeral osteoarthritis.

The medical management and systemic features of RA create some unique challenges in this population. Disease-modifying anti-rheumatic drugs are immunosuppressive and are therefore believed to increase the potential risk of prosthetic joint infection. Corticosteroids are also frequently used to control disease progression and long term use affects bone mineral density increasing the risk of fractures and the quality of implant fixation. Furthermore, the non-orthopaedic

manifestations of the disease increase the risk of systemic peri-operative complications and fitness for anaesthesia.

Historically, anatomical shoulder arthroplasty has played a major role in rheumatoid patients. McCoy *et al.* found good pain relief in up to 93% after surgery [4]. However, despite high patient satisfaction and good pain relief, function is unpredictable and often poor, with patients in some series averaging elevation below the horizontal [5-7]. Barrett *et al.* reported active forward flexion averaged 34 degrees in 140 anatomical shoulder replacements where 50% of patients had significant cuff pathology [8].

Cuff tears in rheumatoid patients are extremely common (20%-100%) [6, 9, 10] and when present, lead to inferior results of joint replacement due to poor pain relief, superior humeral migration, increased shear forces, eccentric wear, instability, early failure and glenoid loosening rates as high as 50% [6, 7, 11-14]. It is therefore clear that the rotator cuff is an important prognosticator of outcome. Patients with intact cuffs get better pain relief and functional improvement [6, 7, 10]. Although successful outcomes after cuff repair at the time of arthroplasty surgery are reported [15] these results should be interpreted with caution as they may not be applicable to a rheumatoid population in whom progressive upward migration of the humeral head (due to failure of the cuff) has been described as inevitable [10]. Furthermore, the morphology of cuff tears in RA patients more frequently involves Infraspinatus, Subscapularis and Teres Minor and this involvement of a larger part of the cuff may also

*Address correspondence to this author at the Southport and Formby District General Hospital, Town Lane, Southport, Merseyside, PR8 6PN, UK; Tel: (+44) 1704 547471; E-mail: edgee@doctors.org.uk

increase the risk of instability and of having a less favourable outcome [3, 16].

Moderate to severe erosive changes of the humeral head and glenoid (Larsen grades 3-5) are seen in up to 68% of rheumatoid patients [17, 18]. Erosion of the glenoid makes implanting a component more challenging, predisposing to less satisfactory fixation, instability and early implant failure. Previous authors advocated hemiarthroplasty as the treatment of choice in patients with large glenoid defects, bypassing issues of replacing the eroded socket [19-21]. However functional results were poor and deteriorated with time after the procedure, predominantly due to progressive glenoid wear [21-23]. Some surgeons still favour hemiarthroplasty in rheumatoid patients with an intact coracoacromial arch, accepting that this may later fail [3, 22]. It should be noted that the use of massive allografts (e.g. tendo-achilles), to stabilize shoulders with rotator cuff and glenoid defects has been tried in conjunction with hemiarthroplasty, with dismal results [24].

RSA offers a potential solution for some of these issues. The biomechanical features of this design confer increased stability (particularly against superior escape) and a significantly improved lever arm for deltoid, potentially resulting in improved range and strength of abduction even in the presence of a deficient rotator cuff [14, 25, 26]. Despite these biomechanical improvements, expert authors have previously cautioned against using RSA in RA patients due to concerns about increased peri-operative complications, poor functional outcomes and high revision rates [27, 28]. The aim of this study is to re-visit the evidence for these concerns by combining data from published series and reporting pooled data on outcomes, complications and technical considerations.

METHODS

Search Strategy

A PubMed search was performed using the terms “reverse shoulder arthroplasty ‘or’ reverse shoulder replacement” and “rheumatoid arthritis ‘or’ inflammatory arthritis” on the 1st of April 2015. Original articles published in English that reported outcomes of RSA in rheumatoid patients were included. Studies were excluded if results were combined with populations containing other aetiologies and the authors were unable to separate out the data.

The PubMed “related citations” tool and Google Scholar were used to search for any further potentially eligible articles. The references of all included articles were also searched for any further relevant articles.

Method of Critical Appraisal

Included studies were appraised by two independent reviewers (EG, EH) with respect to a validated quality assessment scale for retrospective clinical studies developed by Rangel *et al.* [29]. The tool allows identification of the most rigorous evidence by characterising 16 baseline criteria that define thorough and accurate reporting of non-controlled studies. If there was any disagreement between the authors in assigning a score to each paper appraised, a third independent reviewer (AS) made the final decision.

Data Extraction

The main outcomes extracted include: the incidence of complications, the rate of revision, modes of failure and outcome measures relating to function, pain and satisfaction.




Fig. (1). Flow of studies.

Table 1. Adequacy of reporting of studies based on criteria proposed by Rangel *et al.* [29].

Criteria of Rangel <i>et al.</i> [29]	Rittmeister [3]	Hattrup [16]	Ekelund [30]	Holcomb [31]	John [32]	Woodruff [33]	Young [34]
<i>Description of participating surgeons/institutions:</i>							
Can number of centres be determined?	No	✓	No	No	✓	✓	No
Can practice type be determined?	✓	✓	✓	✓	✓	✓	✓
Can number of surgeons be determined?	✓	✓	✓	No	No	✓	✓
Can author's prior experience with procedure be determined?	No	No	No	No	No	No	No
Is timeline of when cases performed documented?	No	No	✓	✓	No	✓	✓
<i>Description and definition of cases:</i>							
Was population from which cases selected described?	✓	✓	✓	✓	✓	No	✓
Are diagnostic criteria clearly documented?	✓	No	No	✓	✓	No	✓
Are eligibility criteria clearly documented?	✓	No	No	✓	✓	No	✓
<i>Description of the intervention:</i>							
Is surgical technique adequately described?	✓	✓	✓	No	✓	✓	✓
Any attempt to standardise operative technique mentioned?	✓	No	No	✓	✓	✓	✓
Any attempt to standardise peri-operative care mentioned?	✓	No	✓	✓	✓	✓	✓
<i>Analysis of outcome data:</i>							
Is mean and range of demographic data reported?	No	✓	✓	No	No	✓	✓
Are outcomes presented with appropriate measures of variability?	✓	✓	✓	✓	✓	✓	✓
Are diagnostic methods for defining outcome described?	✓	✓	✓	✓	✓	✓	✓
Do authors address whether there is missing data?	✓	✓	✓	✓	✓	✓	✓
Is number and nature of complications addressed?	✓	✓	No	✓	No	✓	✓
Total number of criteria satisfied	12/16	10/16	10/16	11/16	11/16	12/16	14/16

RESULTS

After application of the search strategy and eligibility criteria seven studies were included for critical appraisal and data extraction [3, 16, 30-34]. The flow of studies is shown in Fig. (1).

The quality assessment of included studies is presented in Table 1. This shows that included studies were of similar quality and that although there were some weaknesses, there were no major limitations in key areas. Data from these studies were collated to produce an overall study population for this review. This is summarised in Table 2 and comprises a total of 144 shoulders of which 129 had rotator cuff tears identified on pre-operative imaging. After loss to follow up, 121 reverse arthroplasty shoulders were included with weighted mean follow up of 46.9 months (11.5-143 months). The weighted mean age of patients was 64.6 (range 34-86).

Patient-Based Outcome Measures

No single outcome measure was universally reported but all authors reported a significant improvement in the main outcome measure used in their studies. Due to the narrow eligibility criteria used in this review, as evidenced by the

Table 2. Summarises basic characteristics of included studies.

Study	Year	Level of Evidence*	No. of Shoulders	Mean Follow-Up (Months)
Rittmeister [3]	2001	IV	6	54.3 (48-73)
Hattrup [16]	2012	IV	19	37 (24-66)
Ekelund [30]	2011	IV	27	56 (18-143)
Holcomb [31]	2010	IV	21	36 (24-73)
John [32]	2010	III	17	24.3 (11.5-40.9)
Woodruff [33]	2003	IV	13	87 (60-110)
Young [34]	2011	IV	18	46 (25-84)

*Oxford Centre for Evidence-Based Medicine 2011 Levels of Evidence.

small number of included patients, it was considered that the included populations were comparable and that some data could be pooled for further analysis. This is evidenced by the fact that the pre- and post- operative scores for the main outcome measures reported were remarkably similar between studies. This data is presented in Table 3.

The Constant and Murley Score (CMS) evaluates pain, activities of daily living, mobility and strength and was

Table 3. Summary of main outcomes data from included studies.

Study	CS			ASES			Other Objective Outcome Measure	Patient Satisfaction
	Pre-Op	Post-Op	Mean Difference	Pre-Op	Post-Op	Mean Difference		
Rittmeister [3]	17	63	46	-	-	-	-	Overall: 100%, n=6
Hattrup [16]	-	-	-	27	76	49	VAPS: 6.5 - 1 SST: 1 - 8	-
Ekelund [30]	13	52	39	-	-	-	VAPS: 8 - 1	-
Holcomb [31]	-	-	-	28	82	54	VAPS: 7 - 1 SST: 1 - 7	95%(20) Satisfied 5% (1) dissatisfied
John [32]	19	59.5	40.5	-	-	-	SF-36 MSC: 108% SF-36 PSC: 77%	Overall: 93%, n=14
Woodruff [33]	-	58.5	-	-	-	-	SF-12 PSC 38.2 SF-12 MSC 49.13	
Young [34]	22.5	64.9	42.4	-	-	-	MSSS: 68.6%	61%(11) very satisfied 33% (6) satisfied 5% (1) dissatisfied
Weighted mean			40.9			51.6		

Pre-op - pre-operative, *Post-op* - post-operative, *CS* - Constant-Murley Score, *ASES* - American Shoulder and Elbow Surgeons outcome measure, *VAPS* - Visual Analog Pain Scale, *SST* - Simple Shoulder Test, *SF-36 MSC* - Short Form 36 Mental Score Component, *SF-36 PSC* - Short Form 36 Physical Score Component, *SF-12 PSC* - Short Form 12 Physical Score Component, *SF-12 MSC* - Short Form 12 Mental Score Component, *MSSS* - Mean subjective shoulder score.

reported by four studies [3, 30, 32, 34]. The pooled data from these studies included 68 shoulders and the overall weighted mean improvement in the CMS of 40.9 points was seen from 17.4 to 58.3 at an average follow up of 45.3 months for this group of patients.

It is of note that Young *et al.* separated the patients according to the presence of a normal or atrophied Teres Minor as seen on MRI scan. The mean CMS improved from 28.0 points to 74.3 points in 8/15 shoulders with a normal Teres Minor muscle and from 20.8 to 54.6 points in 7/15 shoulders with an atrophic Teres Minor muscle. The seven shoulders with a normal Teres Minor muscle preoperatively all had a final CMS of >70 points, whereas only two of the eight shoulders with Teres Minor atrophy had a final score in this range [34].

Another two studies [16, 31] reported the American Shoulder and Elbow Surgeons (ASES) outcome measure, which evaluates pain and the level of function or disability. These studies included 40 patients and found a weighted mean improvement of 51.6 points from 27.5 to 79.1.

Patient Satisfaction

As well as functional outcome, four studies specifically asked patients to grade overall satisfaction [3, 31, 32, 34]. Of the 60 patients in these studies, 95% (n=57) rated their shoulders as excellent to satisfactory, whereas 5% (n=3) felt the result of their surgery was unsatisfactory. Specific details are not given regarding patients with an unsatisfactory result so causation cannot be identified. Holcomb *et al.* do, however, give individual outcome measures and the patient with an unsatisfactory result reported no pain, an improvement in ASES score from 20 to 82 and was able to abduct and forward flex to 90 degrees post-operatively [31].

Range of Motion

Four of the included studies reported range of movement data pre and post-operatively [16, 30, 31, 34]. The results are presented in Table 4. However, it was not possible to combine data due to different methods of measuring the range and the results being presented in different units. Despite that, in all four studies, the mean forward flexion improved from less than 80° to at least 115°. Abduction was seen to improve in the three studies that reported on it, from less than 70° to more than 100° [16, 30, 31].

Complications

Overall post-operative complications were seen in 19% (n=23) of the 121 shoulders. The incidence and type of complication is summarised in Table 5. Two patients had more than one complication. In one, a deep infection occurred after failed acromial osteosynthesis, post osteotomy, utilised in a trans-acromial approach [3] In the other, fractures of the greater tuberosity and the scapular spine were seen in a single patient post-operatively [34].

Glenoid Bone Loss and the Requirement for Bone Grafting

Of the seven included series, four [16, 30, 31, 34] reported on glenoid bone loss visualised on pre-operative imaging. This was identified in 39 of 85 patients (45%), but only a small number required bone grafting (Ekelund *et al.* n=4/27, Holcomb *et al.* n=5/21, Young 8/18).

Ekelund *et al.* assessed glenoid loss on plain film radiographs for 18 patients and found that eight of the 18 patients assessed had glenoid erosion. Three patients received structural bone grafting and one received cancellous

Table 4. Mean values for range of movement in operated shoulder. Pre-operative, post-operative and the change in range. Rotation values in Ekelund *et al.*'s paper are given in Constant Murley Score (CMS) point form.

	Hatrup [16]			Ekelund [30]			Holcomb [31]			Young [34]		
	Pre-Op	Post-Op	Change	Pre-Op	Post-Op	Change	Pre-Op	Post-Op	Change	Pre-Op	Post-Op	Change
FF	68°	138°	70°	33°	115°	82°	52°	126°	74°	78°	139°	61°
ABD	66°	134°	68°	26°	103°	77°	55°	116°	61°	-	-	-
ER	23°	52°	29°	0.6	5.8	-	19°	33°	14°	15°	20°	5°
IR	-	-	-	2.1	2.9	-	S1	L4	-	sacrum	L3	-

FF - Forward Flexion, ABD - Abduction, ER - External Rotation, IR - Internal Rotation.

graft. One patient also required augmentation of a defect with cement to achieve stability of an inferior screw [30].

Table 5. Summary of Post-operative complications from all papers [3, 16, 30-34].

Post-Operative Complication	Overall Incidence
Infection and septic loosening	3.3%, n= 4
Scapular Spine fracture	3.3%, n= 4
Glenoid fracture	3.3%, n= 4
Failed Acromion Osteosynthesis	2.5%, n= 3
Symptomatic aseptic glenoid loosening	1.7%, n= 2
Greater tuberosity fracture	1.7%, n= 2
Acromial fracture	1.7%, n= 2
Nerve Palsy	1.7%, n= 2
Dislocation	0.8%, n= 1
Coracoid fracture	0.8%, n= 1
Symptomatic aseptic humeral loosening	0%, n= 0
Total	25 complications (23 shoulders = 19%)

Holcomb *et al.* measured available glenoid bone stock (CT/MRI) from the most medial portion of the glenoid to the medial base of the coracoid process, with a mean of 24.2mm (range 11.2-30.5) remaining. The need for grafting wasn't based on these values but on intra-operative visual inspection, if less than 80% of the baseplate had osseous contact, bone grafting (humeral head autograft) was used. Stable fixation of the baseplate was achieved in each case with no cases of non-union of the bone graft [31].

Young *et al.* also assessed glenoid loss using MRI or CT imaging according to Levigne's classification. Fourteen patients (78%) had glenoid erosion. Eight of these were treated intra-operatively with bone grafting, four using cancellous bone autograft and four requiring structural autografting, from the resected humeral head [34] At final follow up, all patients that had received bone grafts had a well fixed glenoid component on plain film radiography.

Hatrup *et al.* took a different approach to managing glenoid erosion and did not use structural bone grafts. Instead, the inferior margin of the glenoid baseplate was aligned with the inferior margin of the glenoid and reaming

was directed to create neutral orientation and over 50% bony contact with the baseplate. No attempt was made to create an inferior tilt for the glenosphere to avoid exacerbating superior bone loss. Cancellous autograft from the humeral head was used to fill residual glenoid defects [16].

Scapula Notching

Notching is a well-recognised consequence of RSA, but the clinical significance is unclear [35, 36] Only four of the included studies reported scapula notching, at a rate of 42%(n=34/81) [16, 30, 32, 34].

The majority of these (21) were grade 1-2 according to the Nerot-Sirveaux classification (appendix) [37]. Thirteen patients had notching above the inferior glenoid component screw. All notching was asymptomatic with no glenoid loosening or fracture being reported as associated with notching within the study periods.

Infection

The overall rate of deep infection in the combined series was 3.3% (n=4) and all four required further surgery [3, 30, 31]. This is comparable to infection rates for this operation for other aetiologies. One patient from the series reported by Ekelund *et al.* underwent a successful two-stage revision after developing deep infection within the first year. Causative organisms were not reported. The Constant-Murley score at latest follow up was 48 points [30].

In Holcomb's series, one patient developed deep infection (*Pseudomonas aeruginosa*) at six years post-operation, requiring removal of the prostheses and a hemiarthroplasty cement spacer. Another patient with early infection (*Methicillin-sensitive Staph. Aureus*) diagnosed at seven weeks required washout, debridement and exchange of polyethylene liner along with six weeks of antimicrobial therapy. This patient was infection free at final follow up (two years) and rated the outcome as 'excellent' [31].

In the Rittmeister *et al.* series, one patient developed sepsis within the joint after re-operation for non-union of acromial osteosynthesis (post osteotomy). This patient required implant removal [3]. Only Ekelund *et al.* reported their antibiotic prophylaxis regimens (Flucloxacillin, or Clindamycin if penicillin allergic) [30]. Holcomb *et al.* advocated the routine inclusion of Tobramycin into bone cement [31]. No clues as to causation of infection can be drawn from the data, nor do the authors postulate causation.

Loosening

There were no cases of symptomatic humeral loosening in the combined series. Twenty cases of radiographic humeral loosening were reported but none of these had symptoms at final review [16, 33]. Woodruff *et al.* found that 100% of their cases had radiolucencies surrounding the uncemented, hydroxyapatite coated humeral components (Delta III) at a mean follow up of 87 months. These may be attributed to stress shielding rather than true loosening and osteolysis.

There were seven (overall incidence 5.7%) cases of glenoid loosening reported. Once again the majority (five patients) were found in the series reported by Woodruff *et al.* and none of these were symptomatic. One of these occurred in a patient that sustained intra-operative fracture of the glenoid [33]. Of the other two, one still rated the shoulder as good [31], the other resulted in breaking of the inferior screw and tilting of the component which required revision [3].

Fracture

Thirteen (10.7%) fractures occurred associated with RSA in this group of patients. Four patients sustained an intra-operative glenoid fracture during glenoid reaming. Holcomb *et al.* fixed the two intra-operative fractures occurring in their series with cannulated lag-screws, allowing subsequent stable implantation of a glenoid component and excellent post-operative results (ASES 80, forward flexion 125 degrees) [31]. Ekelund *et al.* [30] and Woodruff *et al.* [33] both had one intra-operative fracture, neither paper specifically discusses intra-operative fixation or alteration to post-operative management. However, Woodruff, *et al.* reported that this patient had a very deficient glenoid at the time of surgery and found loosening of the glenoid component and a Constant score of 51 at final follow up.

Five fractures were seen in Young's series, two sustaining post-operative acromial fractures, one resulted in inferior tilt of the fragment; the patient rated her outcome as satisfactory, and reported no pain and had active abduction of 140° at the latest follow-up. The other was in the acromial spine; this healed with conservative management but unfortunately led to reduced mobility and the patient reported pain for one year. Two patients sustained avulsion fractures of the greater tuberosity that were thought to have occurred intra-operatively but were not identified at the time. One of these occurred in the patient with an acromial spine fracture and contributed to the poor outcome. The other healed and reported good outcomes. Another patient sustained a coracoid avulsion that was repaired at the time with sutures and healed [34].

In other studies, three more scapular spine fractures occurred after falls; one healed giving excellent results (forward flexion 160 degrees, ASES 83), one was discovered after bone scan for progressive pain, another did not heal and gave a poor outcome with pain score of 10 and ASES of 7 (not revised at final follow up). One patient sustained a stress fracture of the acromion at eight months and although this didn't heal, the patient could forward flex to 100 degrees and the ASES was 60 [16, 30, 31].

Rates of Revision Surgery and Modes of Failure

Rates of revision surgery in the included studies varied from 0-38% at latest follow up. The overall rate of revision was 7% (n=9) at a mean follow up of 46.9 months. However, if the series of six patients reported by Rittmeister *et al.* [3] is excluded, on the basis that it is not contemporary and represents a significant outlier compared to recent studies, then the mean rate of revision is only 5% (n=6/115) in this population at a mean follow up of 46.5 months. Three cases required revision for septic loosening, 1 for instability, 1 for glenoid fracture and 1 due to breakage of the central screw. The three re-operations (50%) described by Rittmeister *et al.* were all related to failed osteosynthesis of the acromion after osteotomy in the trans-acromial approach, used for better exposure, thus adding further weight to the decision to exclude that study in the analysis of revision surgery.

A systematic review by Zumstein *et al.* of 782 patients receiving RSA for all aetiologies, revealed an overall revision rate of 10.1%, which is higher than the rate we found in RA patients [38].

DISCUSSION

RSA is increasingly advocated as an appropriate option for primary arthroplasty in end-stage RA. Of only seven available series specifically focusing on this indication, five were published in the last four years demonstrating a recent increase in popularity for this treatment option.

This review reports outcomes collated from the seven available studies. The most commonly used outcome measure was the Constant-Murley Score (CMS). This includes functional assessment, range of motion and strength of active abduction as well as the pain score [39, 40]. All studies that reported this outcome measure showed a remarkably consistent improvement at a mean of 41.62 (range 39 to 46) points. This is comparable with studies looking at RSA in cuff tears (mean improvement of 43) [19, 41, 42] or cuff arthropathy (mean improvement of 41.33) [43-45] and suggests that outcome from RSA in RA is predictable.

The American Shoulder and Elbow Surgeons (ASES) outcome measure concentrates more on subjective functional limitations of the shoulder and pain associated with it. Studies included in our review showed an overall mean improvement in ASES scores of 51.6 points (from 27.5 to 79.1), which, although not directly comparable, is greater than the improvement reported in RSA performed for cuff arthropathy (40.75) [46-49].

John *et al.* reported results from the Short-Form 36 (SF-36). Although there was some improvement in physical function, a great improvement was seen in the mental and emotional components, above that of a normative population. The authors postulated this might be due to lower expectations of function in RA patients and an increase in satisfaction once pain had been relieved, despite still having a functional impairment [32]. This theory can also be used to explain high satisfaction rates seen in this lower demand and lower expectation cohort burdened with chronic functional impairment and pain.

Range of movement was seen to improve in all mean values in the four studies in which they were reported [16, 30, 31, 34]. It was not possible to combine data for analysis but there are some key points to note. Firstly, as stated in the introduction forward elevation in RA patients after TSA or hemiarthroplasty can be extremely poor [5-8, 21-23]. Khan compared osteoarthritic patients to RA patients undergoing anatomical total shoulder arthroplasty and found post-operative forward elevation averaging 130° and 85° respectively [6] Table 4 demonstrates that the authors of the studies reporting on range of motion included in this review all recorded a minimum mean forward elevation of 115 degrees. Although a comparative study is required to demonstrate a significant difference in ROM this data certainly reinforces the empirical observation of many surgeons that RSA allows greater potential for forward elevation than anatomical replacement in an RA population due to the high incidence of associated cuff tears.

Furthermore, although Young *et al.* found little improvement in external rotation with the arm in an adducted position, with mean external rotation improving from 15° to 20° of rotation, they also tested 'functional external rotation', with the arm at 90° abduction. They found a greater improvement from 17° pre-operatively to 46° functional external rotation post-operatively. The authors attributed this difference to the presence of a functioning Teres Minor muscle facilitating external rotation with the arm in this position. This also accounted for the difference in CMS scores reported in this paper [34].

Shoulder arthroplasty complication rates have declined in recent years with a lower need for re-operation. It is stated by some authors that there is no discernable link between early or late complications and initial aetiology, with comparable complication rates occurring in RSA regardless of whether the patient has RA [50]. In order to ascertain whether the perceived increase in risk of complication, when performing RSA in RA patients, does exist, comparisons can be made to RSAs performed for mixed aetiologies. A systematic review by Zumstein *et al.* looked at the results of 782 RSAs in 761 patients performed for multiple aetiologies, including cuff tears, fracture, tumour, revision surgery and RA [38].

The overall complication rate in this series of patients was 19% (23/121 shoulders). Wall *et al.* reported a similar complication rate of 19% in a series of 191 reverse arthroplasties [51] and Frankle *et al.* found a complication rate of 17% [52]. Zumstein *et al.* actually found a higher rate of complication in their systematic review of 782 RSAs for all aetiologies. They separated 'problems' and 'complications', based on whether they thought the final outcome was likely (complication) or unlikely (problem) to be affected. They reported an overall 'problem' rate of 44% and a 'complication' rate of 24% in a mixed population [38].

One of the main concerns when pre-operatively planning shoulder arthroplasty in RA patients is glenoid bone loss, and this was apparent in 45% of patients in this cohort. Less than 20% of those with bone loss (9% overall) required grafting. By final follow up, all these patients had a well fixed and stable glenoid component with no signs of non-union or early loosening, suggesting that this is a useful option for managing deficient glenoid bone stock.

The rate of scapula notching that occurred in the combined series was 34%. Prior clinical studies have shown scapular notching rates ranging from 0% to 96% in RSA [53]. Notching is usually apparent within the first few post-operative months and very few shoulders that have avoided notching by 12 months demonstrate notching later [28, 42, 54]. This suggests that the length of follow up in the combined series was of sufficient duration to appropriately capture the incidence of scapula notching and that its incidence was no greater than that seen when RSA was performed for all aetiologies (35%) [38]. No direct links were made between notching and fracture, loosening or revision in any of the studies included in this review.

Infection is a devastating complication in arthroplasty. RA patients are potentially immunocompromised and so higher infection rates are anticipated. The overall infection rate in this combined series was 3.3%, which is not dissimilar to a population of mixed aetiologies undergoing the same operation. In previous papers concerning a normal population undergoing primary RSA, infection rate is stated as between 1% and 5% [36, 55], and Zumstein *et al.*'s review of 782 RSAs found a deep infection rate of 3.8% [38]. The infecting organisms described in the patient cohort included in this review were typical, including *Pseudomonas acnes*, *Pseudomonas aeruginosa*, *Staphylococcus Aureus*, and coagulase-negative *Staphylococcus* [31, 56]. However, rare cases of prosthetic joint infection in RSA implants with *Candida Albicans* have been reported in RA patients. It is hypothesised that this is almost certainly due to immunosuppression caused by disease controlling drugs such as anti-tumour necrosis factor therapy (e.g. Etanercept) and should also therefore be considered as a potential infecting micro-organism if an RA patient presents with a prosthetic joint infection [57].

Osteopenia and osteoporosis associated with glucocorticoid medication, disuse and the disease process itself, raise concerns of fracture in RA patients. Thirteen fractures were seen in twelve patients within the cohort included in this review, giving a fracture risk of 10.7% overall. Seven occurred intra-operatively and six occurred post-operatively. Seven of these fractures healed and in general these reported good outcomes, apart from the patient with an acromial spine and a greater tuberosity fracture who had an unsatisfactory outcome. Five fractures did not heal and four of these reported poor outcomes. The healing and outcome of one of the fractures was not reported. Zumstein *et al.* reported 46 fractures in 782 RSAs in a mixed population, giving an overall fracture risk of 5.9%, which is lower than seen in the purely RA cohort of this review [38]. As the rate of fracture varied greatly between the studies in this review from 4% [30] to 28% [34], it may be that specific aspects of the intra-operative technique or the prosthesis used increased the risk of fracture in certain studies however this cannot be ascertained from the data provided.

There were no cases of symptomatic humeral loosening and only two cases of symptomatic glenoid loosening, one of which required revision in the combined series. Depending on the criteria used for radiographic loosening, extremely high rates of up to 100% were reported and Woodruff *et al.* attributed this to stress shielding surrounding uncemented components rather than true osteolysis [33]. Zumstein *et al.*

found radiographic loosening in 27 (out of 782) RSAs giving an overall rate of 3.5% in a mixed aetiology population. Loosening rates varied greatly depending on which prosthesis was used and the fixation type. The authors found loosening rates were twice as high in the Encore system compared to the Grammont system [38].

A final concern with rheumatoid patients is the existence of multi-system involvement and a perceived increased risk of peri-operative systemic complications. None of the included studies reported systemic peri-operative complications. Hambright *et al.* reported a comparison of peri-operative complications in RA vs non RA patients undergoing shoulder arthroplasty and found this to be less than 1% in both groups, with no significant difference between them [58].

Due to a limited number of studies providing data on this subject and a variation of follow up, we are unable to report cumulative survivorship. However, the mean rate of revision from the contemporaneous combined series (excluding Rittmeister *et al.* [3] as previously discussed) was 5% at a mean follow up of 46.9 months. Although not directly comparable, when one is considering the role of RSA in RA patients it is important to note that the revision rate given for RSA in a mixed population was 10.1% suggesting that the risk for RA patients is not excessively high [38]. Other studies have also noted that revision rates for anatomical shoulder arthroplasty in RA patients are lower than those published for non-RA patients, possibly due to a lower demand in this cohort or to statistical variance secondary to small sample sizes [21, 59-61].

CONCLUSION

The contemporary literature does not show a higher rate of complications for RSA in rheumatoid patients either in the peri- or post-operative period when compared to RSA in a large population of patients with mixed aetiologies [38]. Pre-conceptions of increased rates of infection, fracture, notching, loosening and failure were not realised in this review when compared to other aetiologies.

The data suggests it is a safe procedure that allows predictable improvements in the ROM, CMS and ASES scores as well as high patient satisfaction. Longer-term data, larger numbers and prospective, controlled trials may be beneficial in substantiating this and to better estimate long-term survivorship. This review concludes that RSA is a valuable treatment option for end stage arthritis in rheumatoid patients.

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Cuomo F, Greller MJ, Zuckerman JD. The rheumatoid shoulder. *Rheum Dis Clin North Am* 1998; 24: 67-82.
- [2] Lehtinen JT, Kaarela K, Belt EA, Kautiainen HJ, Kauppi MJ, Lehto MU. Incidence of glenohumeral joint involvement in seropositive rheumatoid arthritis. A 15 year endpoint study. *J Rheumatol* 2000; 27: 347-50.
- [3] Rittmeister M, Kerschbaumer F. Grammont reverse total shoulder arthroplasty in patients with rheumatoid arthritis and nonreconstructible rotator cuff lesions. *J Shoulder Elbow Surg* 2001; 10(1): 17-22.
- [4] McCoy S, Warren RF, Bade HA, Ranawat CS, Inglis AE. Total shoulder arthroplasty in Rheumatoid Arthritis. *J Arthroplasty* 1989; 4: 105-13.
- [5] Jolles BM, Grosso P, Bogoch ER. Shoulder arthroplasty in patients with juvenile rheumatoid arthritis. *J Arthroplasty* 2007; 22: 876-83.
- [6] Khan A, Bunker TD, Kitson JB. Clinical and radiological follow-up of the Aequalis third-generation cemented total shoulder replacement. A minimum ten-year study. *J Bone Joint Surg Br* 2009; 91:1594-1600. *J Bone Joint Surg Br* 2009; 91(12): 1594-600.
- [7] Sperling JW, Cofield RH, Schleck CD, Harmsen WS. Total shoulder arthroplasty versus hemiarthroplasty for rheumatoid arthritis of the shoulder: results of 303 consecutive cases. *J Shoulder Elbow Surg* 2007; 16(6): 683-90.
- [8] Barrett WP, Thornhill TS, Thomas WH, Gerbhart EM, Sledge CB. Non-constrained total shoulder arthroplasty in patients with polyarticular rheumatoid arthritis. *J Arthroplasty* 1989; 4: 91-6.
- [9] Ennevaara K. Painful shoulder joint in rheumatoid arthritis: a clinical and radiographic study of 299 cases with special reference to arthrography of the glenohumeral joint. *Acta Rheumatol Scand* 1967; (Suppl 11): 111-6.
- [10] Lehtinen JT, Belt EA, Lybäck CO, *et al.* Subacromial space in the rheumatoid shoulder: a radiographic 15-year follow-up study of 148 shoulders. *J Shoulder Elbow Surg* 2000; 9: 183-7.
- [11] Cofield RH. Total shoulder arthroplasty with the Neer prosthesis. *J Bone Joint Surg* 1984; 66: 899-906.
- [12] Franklin JL, Barrett WP, Jackins SE, Matsen FA 3rd. Glenoid loosening in total shoulder arthroplasty: association with rotator cuff deficiency. *J Arthroplasty* 1977; 3(1): 39-46.
- [13] Khan WS, Longo UG, Ahrens PM, Denaro V, Maffulli N. A systematic review of the reverse shoulder replacement in rotator cuff arthropathy, rotator cuff tears, and rheumatoid arthritis. *Sports Med Arthrosc* 2011; 19(4): 366-79.
- [14] Smithers CJ, Young AA, Walch G. Reverse shoulder arthroplasty. *Curr Rev Musculoskelet Med.* 2011; 4(4): 183-90.
- [15] Rozing PM, Brand R. Rotator cuff repair during shoulder arthroplasty in rheumatoid arthritis. *J Arthroplasty* 1998; 13: 311-9.
- [16] Hattrup SJ, Sanchez-Sotelo J, Sperling JW, Cofield RH. Reverse shoulder replacement for patients with inflammatory arthritis. *J Hand Surg* 2012; 37(9): 1888-94.
- [17] Larsen A, Dale K, Eek M. Radiographic evaluation of rheumatoid arthritis and related conditions by standard reference films. *Acta Radiol Diagn (Stockh)* 1977; 18(4): 481-91.
- [18] Tanaka H, Sugamoto K, Sahara W, *et al.* The mode of destruction in shoulders with rheumatoid arthritis based on radiographic findings. *J Shoulder Elbow Surg* 2007; 16(6): 539-43.
- [19] Grassi FA, Murena L, Valli F, Alberio R. Six-year experience with Delta III reverse shoulder prosthesis. *J Orthop Surg (Hong Kong)* 2009; 17(2): 151-6.
- [20] Sojbjerg JO, Frich LH, Johannsen HV, Sneppen O. Late results of total shoulder replacement in patients with rheumatoid arthritis. *Clin Orthop Relat Res* 1999; 366: 39-45.
- [21] Van de Sande MA, Brand R, Rozing PM. Indications, complications, and results of shoulder arthroplasty. *Scand J Rheumatol* 2006; 35(6): 426-34.
- [22] Boyer P, Hutten D, Alnot JY. Bipolar shoulder prosthesis for rheumatoid arthritis with irreparable rotator cuff tear: results after 5 years' follow up [Article in French]. *Rev Chir Orthop Reparatrice Appar Mot* 2006; 92(6): 556-66.
- [23] Gregory T, Hansen U, Emery RJ, Augereau B, Amis AA. Development in shoulder arthroplasty. *Proc Inst Mech Eng H* 2007; 221(1): 87-96.

- [24] Norris TR, Kelly JDI, Humphrey CS. Management of glenoid bone defects in revision shoulder arthroplasty: a new application of the reverse total shoulder prosthesis. *Tech Shoulder Elbow Surg* 2007; 8: 37-46.
- [25] Grammont PM, Baulot E. Delta shoulder prosthesis: biomechanical principles. [Personal communication] 1991.
- [26] Grammont PM, Baulot E. Delta shoulder prosthesis for rotator cuff rupture. *Orthopedics* 1993; 16: 65-8.
- [27] Gerber C, Pennington SD, Nyffeler RW. Reverse total shoulder arthroplasty. *J Am Acad Orthop Surg* 2009; 17(5): 284-95.
- [28] Guery J, Favard L, Sirveaux F, Oudet D, Mole D, Walch G. Reverse total shoulder arthroplasty. Survivorship analysis of eighty replacements followed for five to ten years. *J Bone Joint Surg Am* 2006; 88(8): 1742-7.
- [29] Rangel S, Kelsey J, Henry MCW, Moss RL. Critical analysis of clinical research reporting in pediatric surgery: Justifying the need for a new standard. *J Pediatr Surg* 2003; 38(12): 1739-43.
- [30] Ekelund A, Nyberg R. Can Reverse Shoulder Arthroplasty be Used with Few Complications in Rheumatoid Arthritis? *Clin Orthop Relat Res* 2011; 469(9): 2483-8.
- [31] Holcomb JO, Hebert DJ, Mighell MA, et al. Reverse shoulder arthroplasty in patients with rheumatoid arthritis. *J Shoulder Elbow Surg* 2010; 19(7): 1076-84.
- [32] John M, Pap G, Angst F, et al. Short-term results after reversed shoulder arthroplasty (Delta III) in patients with rheumatoid arthritis and irreparable rotator cuff tear. *Int Orthop* 2010; 34(1): 71-7.
- [33] Woodruff MJ, Cohen AP, Bradley JG. Arthroplasty of the shoulder in rheumatoid arthritis with rotator cuff dysfunction. *Int Orthop* 2003; 27(1): 7-10.
- [34] Young A, Smith M, Bacle G, Moraga C, Walch G. Early results of reverse shoulder arthroplasty in patients with rheumatoid arthritis. *J Bone Joint Surg Am* 2011; 93: 1915-23.
- [35] Sirveaux F, Favard L, Oudet, Huquet D, Walch G, Molé D. Grammont inverted total shoulder arthroplasty in the treatment of glenohumeral osteoarthritis with massive rupture of the cuff. *J Bone Joint Surg Br* 2004; 86: 388-95.
- [36] Guery J, Favard L, Sirveaux F, Oudet D, Molé D, Walch G. Reverse total shoulder arthroplasty. Survivorship analysis of eighty replacements followed for five to ten years. *J Bone Joint Surg Am* 2006; 88: 1742-7.
- [37] Sirveaux F. La prothèse de Grammont dans le traitement des arthropathies de l'épaule à coiffe détruite. A propos d'une série multicentrique de 42 cas. Faculté de médecine de Nancy, thèse de l'université de Nancy I 1997; 245.
- [38] Zumstein MA, Pinedo M, Old J, Boileau P. Problems, complications, reoperations, and revisions in reverse total shoulder arthroplasty: A systematic review. *J Shoulder Elbow Surg* 2011; 20: 146-57.
- [39] Boehm D, Wollmerstedt N, Doesch M, Handwerker M, Mehling E, Gohlke F. Development of a questionnaire based on the Constant-Murley-Score for self-evaluation of shoulder function by patients. [Article in German]. *Unfallchirurg* 2004; 107(5): 397-402.
- [40] Conboy VB, Morris RW, Kiss J, Carr AJ. An evaluation of the Constant-Murley shoulder assessment. *J Bone Joint Surg Br* 1996; 78(2): 229-32.
- [41] Boileau P, Gonzalez JF, Chuinard C, Bicknell R, Walch G. Reverse total shoulder arthroplasty after failed rotator cuff surgery. *J Shoulder Elbow Surg* 2009; 18(4): 600-6.
- [42] Sirveaux F, Favard L, Oudet D, Huquet D, Walch G, Molé D. Grammont inverted total shoulder arthroplasty in the treatment of glenohumeral osteoarthritis with massive rupture of the cuff Results of a Multicentre Study of 80 Shoulders. *J Bone Joint Surg Br* 2004; 86(3): 388-95.
- [43] Boileau P, Watkinson D, Hatzidakis AM, Hovorka I. Neer award 2005: the Grammont reverse shoulder prosthesis: results in cuff tear arthritis, fracture sequelae, and revision arthroplasty. *J Shoulder Elbow Surg* 2006; 15: 527-40.
- [44] Neyton L, Boileau P, Nové-Josserand L, Edwards TB, Walch G. Glenoid bone grafting with a reverse design prosthesis. *J Shoulder Elbow Surg* 2007; 16(Suppl 3): 71-8.
- [45] Sayana MK, Kakarala G, Bandi S, Wynn-Jones C. Medium term results of reverse total shoulder replacement in patients with rotator cuff arthropathy. *Ir J Med Sci* 2009; 178(2): 147-50.
- [46] Cuff D, Pupello D, Virani N, Levy J, Frankle M. Reverse shoulder arthroplasty for the treatment of rotator cuff deficiency. *J Bone Joint Surg Am* 2008; 90: 1244-51.
- [47] Klein SM, Dunning P, Mulieri P, Pupello D, Downes K, Frankle MA. Effects of acquired glenoid bone defects on surgical technique and clinical outcomes on reverse shoulder arthroplasty. *J Bone Joint Surg Am* 2010; 92: 1144-54.
- [48] Mullieri P, Dunning P, Klein S, Pupello D, Frankle M. Reverse shoulder arthroplasty for the treatment of irreparable rotator cuff tear without glenohumeral arthritis. *J Bone Joint Surg Am* 2010; 92: 2544-56.
- [49] Werner CM, Steinmann PA, Gilbert M, Gerber C. Treatment of painful pseudoparesis due to irreparable rotator cuff dysfunction with the Delta III reverse-ball-and-socket total shoulder prosthesis. *J Bone Joint Surg Am* 2005; 87: 1476-86.
- [50] Chin PY, Sperling JW, Cofield RH, Schleck C. Complications of total shoulder arthroplasty: are they fewer or different? *J Shoulder Elbow Surg* 2006; 15: 19-22.
- [51] Wall B, Nové-Josserand L, O'Connor LP, Edwards TB, Walch G. Reverse total shoulder arthroplasty: a review of results according to etiology. *J Bone Joint Surg Am* 2007; 89: 1476-85.
- [52] Frankle M, Siegal S, Pupello D, Saleem A, Mighell M, Vasey M. The reverse shoulder prosthesis for glenohumeral arthritis associated with severe rotator cuff deficiency. A minimum two-year follow-up study of sixty patients. *J Bone Joint Surg Am* 2005; 87: 1697-705.
- [53] Kempton LB, Balasubramaniam M, Ankerson E, Wiater JM. A radiographic analysis of the effects of glenosphere position on scapular notching following reverse total shoulder arthroplasty. *J Shoulder Elbow Surg* 2011; 20: 968-74.
- [54] Nyffeler RW, Werner CM, Simmen BR, Gerber C. Analysis of a retrieved Delta III total shoulder prosthesis. *J Bone Joint Surg Br* 2004; 86: 1187-91.
- [55] Trappey GJ, O'Connor DP, Bradley Edwards T. What are the instability and infection rates after reverse shoulder arthroplasty? *Clin Orthop Relat Res* 2011; 469: 2505-11.
- [56] Kelly JD II, Zhao JX, Rhett Hobgood E, Norris TR. Clinical results of revision shoulder arthroplasty using the reverse prosthesis. *J Shoulder Elbow Surg* 2012; 21: 1516-20.
- [57] Radha S, Siddiqui BM, Senevirathna S, Harrison JW. Candida albicans infection of a reverse-polarity shoulder replacement in a patient with rheumatoid arthritis treated with etanercept. *J Clin Rheumatol* 2012; 18(6): 323.
- [58] Hambright D, Henderson RA, Cook C, Worrell T, Moorman CT, Bolognesi MP. A comparison of perioperative outcomes in patients with and without rheumatoid arthritis after receiving a total shoulder replacement arthroplasty. *J Shoulder Elbow Surg* 2011; 20(1): 77-85.
- [59] Brenner BC, Ferlic DC, Clayton ML, Dennis DA. Survivorship of unconstrained total shoulder arthroplasty. *J Bone Joint Surg Am* 1989; 71: 1289-96.
- [60] Neer CS II, Kirby RM. Revision of humeral head and total shoulder arthroplasties. *Clin Orthop Relat Res* 1982; 189-95.
- [61] Wirth MA, Seltzer DG, Senes HR, Pannone A, Lee AJ, Rockwood CA. An analysis of failed humeral head and total shoulder arthroplasty [abstract]. *J Shoulder Elbow Surg* 1995; 4(Suppl 1): S13.